

Bilateral Fixed-Points and Algebraic Properties of Viability Kernels and Capture Basins of Sets

Jean-Pierre Aubin & Francine Catté

Université Paris-Dauphine

F-75775 Paris cx (16)

e-mail: J.P.Aubin@wanadoo.fr, Catte@ceremade.dauphine.fr

Mars 23, 2001

Abstract

Many concepts of viability theory such as viability or invariance kernels and capture or absorption basins under discrete multivalued systems, differential inclusions and dynamical games share algebraic properties that provide simple — yet powerful — characterizations as either largest or smallest fixed points or unique minimax (or bilateral fixed-point) of adequate maps defined on pairs of subsets. Further, important algorithms such as the Saint-Pierre viability kernel algorithm for computing viability kernels under discrete system and the Cardaliaguet algorithm for characterizing “discriminating kernels” under dynamical games are algebraic in nature. The Matheron Theorem as well as the Galois transform find applications in the field of control and dynamical games allowing us to clarify concepts and simplify proofs.

keywords viability kernel, capture basin, discriminating kernel, Matheron Theorem, Saint-Pierre viability kernel algorithm, Cardaliaguet discriminating kernel algorithm, openings, closings, Galois transform.

AMS Classification: 93C10, 93C15, 93C55, 49J24, 49J40, 49J53

1 Introduction

Let us consider a differential inclusion $x' \in F(x)$ (summarizing the dynamics of a control system) and two subsets C and K of a finite dimensional vector space X such that $C \subset K$. Here, K is regarded as a constrained subset in which the solution must evolve until possibly reaching the subset C regarded as a target. We recall the following definitions:

1. The subset $\text{Viab}_F(K, C)$ of initial states $x_0 \in K$ such that at least one solution $x(\cdot)$ to differential inclusion $x' \in F(x)$ starting at x_0 is viable in K for all $t \geq 0$ or viable in K

until it reaches C in finite time is called the viability kernel of K with target C under F . When the target $C := \emptyset$ is empty, we say that $\text{Viab}(K) = \text{Viab}(K, \emptyset)$ is the viability kernel $\text{Viab}(K)$ of K .

2. The subset $\text{Capt}_F(K, C)$ of initial states $x_0 \in K$ such that C is reached in finite time before possibly leaving K by at least one solution $x(\cdot)$ to differential inclusion $x' \in F(x)$ starting at x_0 is called the viable-capture basin of C in K and $\text{Capt}_F(C) := \text{Capt}_F(X, C)$ is said to be the capture basin of C .

The subset $\text{Env}_F(C) := \text{Capt}_{-F}(C)$ is known under various names such as invariance envelope or accessibility map or controlled map of C (See [57, Quincampoix] for properties of invariance envelopes under Lipschitz maps and [6, Aubin] for Marchaud maps). Henri Poincaré introduced the concept of shadow (in French, ombre) of K , which is the set of initial points of K from which (all) solutions leave K in finite time. It is thus equal to the complement $K \setminus \text{Viab}_F(K)$ of the viability kernel of K , which has been introduced in the context of differential inclusions in [1, Aubin]. The concept of viability kernel with a target by a Lipschitz set-valued map has been introduced and studied in [60, Quincampoix & Veliov] and the viable-capture basin in [10, Aubin].

Many applications of these concepts have been obtained (see [2, 3, Aubin], [34, Cardaliaguet, Quincampoix & Saint-Pierre] for a presentation of some of them). Applied to subsets such as epigraphs of extended real-valued functions or graphs of maps, they provide interesting existence and uniqueness results of adequately generalized solutions to systems of Hamilton-Jacobi inclusions (see [18, Aubin & Frankowska], [10, 7, Aubin] and, in mathematical finance, [56, Pujal], [23, Aubin, Pujal & Saint-Pierre]).

Various characterizations of the viability kernel and the capture basins finding their origin in the papers of Frankowska on Hamilton-Jacobi equations (see for instance [42, 43, Frankowska]) have been proposed in [10, Aubin] in terms of viability and backward invariance properties and thus, thanks to the Viability and Invariance Theorems, in terms of tangential or normal conditions.

Actually, fixed point properties of the two maps $K \mapsto \text{Viab}_F(K, C)$ and $C \mapsto \text{Viab}_F(K, C)$ popped up here and there, in the study of such and such application, until it was observed that the viability kernel $\text{Viab}_F(K, C)$ of K with target C is the unique “minimax (bilateral fixed-point)” D between C and K of the map $(K, C) \mapsto \text{Viab}_F(K, C)$ in the sense that

$$C \subset D \subset K \ \& \ D = \text{Viab}(K, D) = \text{Viab}(D, C)$$

In other words, saying that a subset K is viable outside C if it coincides with its viability kernel K with target C :

$$K = \text{Viab}_F(K, C)$$

and that a subset $C \subset K$ is isolated in K by F if it coincides with the viability kernel K with target C :

$$\text{Viab}_F(K, C) = C$$

we can restate the above result by stating that the viability kernel $\text{Viab}_F(K, C)$ of K with target C is the unique subset D viable outside C and isolated in K :

$$\text{Viab}(K, C) = \text{Viab}(\text{Viab}(K, C), C) = \text{Viab}(K, \text{Viab}(K, C))$$

This is indeed the natural starting point for further characterizations of the above two fixed-point properties, for instance in terms of local viability of $K \setminus C$ and backward invariance of C when viability kernels are concerned, as in [10, Aubin].

These algebraic properties are valid for other concepts that had been introduced in viability theory, such as invariance kernels $\text{Inv}(K, C)$ of K with target C and invariance-absorption basins $\text{Abs}(K, C)$ under differential inclusions as we shall see in the following pages.

They are valid not only for differential inclusions, but also for differential inclusions with memory and for general evolutionary systems $\mathcal{S} : X \rightsquigarrow \mathcal{C}(0, \infty; X)$ defined later in the paper, as well as for conditional and guaranteed viability kernels and capture basins in dynamical games (see [27, 28, 30, 31, Cardaliaguet]) and evolutionary games. They also hold true for impulse differential inclusions and hybrid systems, as it is shown in [19, Aubin & Haddad] (for more details on hybrid systems in the framework of viability theory, see [5, 9, 8, Aubin], [20, 21, Aubin & Haddad], [22, Aubin, Lygeros, Quincampoix, Sastry & Seube], [35, 36, Cruck], [62, Saint-Pierre]), qualitative viability kernel in [12, Aubin & Dordan], that are not presented in this paper.

Two words about this algebraic framework, that is also closely related to algebraic properties of mathematical morphology (see [52, Matheron], [63, 64, Serra], [45, Giardina & Dougherty], [55, Schmitt & Mattioli], [50, Heijmans] for instance) and max-plus algebras (see [24, Bacelli, Cohen, Olsder & Quadrat] and [44, Gaubert]). Indeed, they rely only on the following two properties of maps $(K, C) \mapsto \mathcal{A}(K, C)$:

$$\left\{ \begin{array}{l} i) \quad C \subset \mathcal{A}(K, C) \subset K \text{ (extensivity/antiextensivity)} \\ ii) \quad (K, C) \mapsto \mathcal{A}(K, C) \text{ is increasing} \end{array} \right.$$

shared by the above viability kernels and capture basins. This has already been noticed first in [54, Mattioli, Doyen & Najman] and then, taken up in chapter 7 of [4, Aubin] for the partial maps $K \mapsto \text{Viab}(K)$ before the systematic study of viability kernels $\text{Viab}(K, C)$ with targets and capture basins $\text{Capt}(K, C)$ of targets viable in a constrained subset that involves functions defined on pairs of subsets.

This “minimax (bilateral fixed-point)” characterization, when valid, underlies some properties of [7, 11, Aubin], [56, Pujal], [23, Aubin, Pujal & Saint-Pierre]. In the same way,

the Galois transform of the invariance kernel map $K \mapsto \text{Inv}_F(K, C)$ provides the invariance envelope.

Not all maps $(K, C) \mapsto \mathcal{A}(K, C)$ defined on the family of subsets do enjoy this minimax property as the viability kernel map does, for instance. In this case, Matheron's theorem allows us to associate with \mathcal{A} maps $K \mapsto \mathcal{A}^\sharp(K, C)$ and $C \mapsto \mathcal{A}^\flat(K, C)$ that have fixed points. They find applications here by providing alternate characterizations of the viability kernel under discrete systems or discriminating kernels under dynamical games. They project a new light on the viability kernel and capture basin algorithms introduced in [61, Saint-Pierre] and the Cardaliaguet algorithm for constructing discriminating kernels under dynamical games in [27, 28, 30, 31, Cardaliaguet] and [34, Cardaliaguet, Quincampoix & Saint-Pierre].

Although the abstract algebraic results we present below are valid on any lattice, we present them only on sub-lattices of the family of subsets of a given set X ordered with the usual inclusion.

It usually takes a long time to discover the sources of a river — and to decide which is the real source. By now, one could “presume” that a source to uniqueness theorems of solutions to Hamilton-Jacobi-Bellman equations as the ones of [40, 41, 42, 43, Frankowska]) based on backward viability and invariance of epigraphs lies with these simple algebraic results that we now describe, and it took that time to isolate this source, indeed.

2 Minimax (bilateral fixed-points) of an opening/closing map

This section is devoted to the existence and uniqueness of maps $(K, C) \mapsto \mathcal{A}(K, C)$ that are pre-openings with respect to K and opening with respect to C .

2.1 Openings and closings

Definition 2.1 *Let us consider a family $\mathcal{D} \subset \mathcal{P}(X) := 2^X$ of subsets of a space X stable under the operations of union and intersection.*

We associate with any pair $(K, C) \in \mathcal{D} \times \mathcal{D}$ the family

$$\mathcal{D}(K, C) := \{D \in \mathcal{D} \mid C \subset D \subset K\}$$

Let us consider a map $\mathcal{A} : \mathcal{D} \mapsto \mathcal{D}$, We shall say that

1. *the map $K \mapsto \mathcal{A}(K, C)$ is a pre-opening if it is increasing (in the sense that whenever $K_1 \subset K_2$, then $\mathcal{A}(K_1, C) \subset \mathcal{A}(K_2, C)$) and antiextensive (in the sense that for any pair $(K, C) \in \mathcal{D}$, $\mathcal{A}(K, C) \subset K$),*
2. *the map $C \mapsto \mathcal{A}(K, C)$ is a pre-closing if it is increasing (in the sense that whenever*

$C_1 \subset C_2$, then $\mathcal{A}(K, C_1) \subset \mathcal{A}(K, C_2)$) and extensive (in the sense that for any pair $(K, C) \in \mathcal{D}$, $C \subset \mathcal{A}(K, C)$),

We recall that the map $D \mapsto \mathcal{A}(D, C)$ is an opening if it is a pre-opening and an idempotent map in the sense that

$$\forall (K, C) \in \mathcal{D}^2, \mathcal{A}(\mathcal{A}(K, C), C) = \mathcal{A}(K, C)$$

and that the map $D \mapsto \mathcal{A}(K, D)$ is a closing if it is a pre-closing and an idempotent map in the sense that

$$\forall (K, C) \in \mathcal{D}^2, \mathcal{A}(K, \mathcal{A}(K, C)) = \mathcal{A}(K, C)$$

We observe that

- $\mathcal{D}(K, \emptyset)$ is the family of subset $D \in \mathcal{D}$ contained in K ,
- $\mathcal{D}(X, C)$ is the family of subset $D \in \mathcal{D}$ containing C .

As we see, these properties involve only the order relations on the family $\mathcal{P}(X)$ of subsets of X , or on any sub-family $\mathcal{D} \subset \mathcal{P}(X)$ stable under the union (“max-closed”) or intersection operations (“min-closed”). Although they can be presented for arbitrary lattices or semi-lattices, we present them in the familiar context of families of subsets.

We next associate with \mathcal{A} the two following family of subsets:

1. the family $\mathcal{K}_{\mathcal{A}}(K, C)$ defined by

$$\mathcal{K}_{\mathcal{A}}(K, C) := \{D \in \mathcal{D}(K, C) \mid D \subset \mathcal{A}(D, C)\}$$

2. the family $\mathcal{I}_{\mathcal{A}}(K, C)$ defined by

$$\mathcal{I}_{\mathcal{A}}(K, C) := \{D \in \mathcal{D}(K, C) \mid \mathcal{A}(K, D) \subset D\}$$

We check at once the following claim:

Lemma 2.2 *If the map $D \mapsto \mathcal{A}(D, C)$ is antiextensive, then $\mathcal{K}_{\mathcal{A}}(K, C)$ is the set of fixed-points of the map $D \mapsto \mathcal{A}(D, C)$ between C and K :*

$$\mathcal{K}_{\mathcal{A}}(K, C) := \{D \in \mathcal{D}(K, C) \mid D = \mathcal{A}(D, C)\}$$

Symmetrically, if the map $D \mapsto \mathcal{A}(K, D)$ is extensive, then $\mathcal{I}_{\mathcal{A}}(K, C)$ is the set of fixed-points of the map $D \mapsto \mathcal{A}(K, D)$ between C and K :

$$\mathcal{I}_{\mathcal{A}}(K, C) := \{D \in \mathcal{D}(K, C) \mid \mathcal{A}(K, D) = D\}$$

Furthermore,

1. if the map $D \mapsto \mathcal{A}(D, C)$ is a pre-opening, then

$$\forall D \in \mathcal{K}_{\mathcal{A}}(K, C), \quad D \subset \mathcal{A}(K, C)$$

2. if the map $D \mapsto \mathcal{A}(K, D)$ is a pre-closing, then for any

$$\forall D \in \mathcal{I}_{\mathcal{A}}(K, C), \quad \mathcal{A}(K, C) \subset D$$

Consequently, if the map $(K, C) \mapsto \mathcal{A}(K, C)$ is a pre-opening with respect to K and a pre-closing with respect to C , then C belongs to $\mathcal{K}_{\mathcal{A}}(K, C)$, K belongs to $\mathcal{I}_{\mathcal{A}}(K, C)$ and

$$\forall D \in \mathcal{D}(K, C), \quad C \subset \bigcup_{D \in \mathcal{K}_{\mathcal{A}}(K, C)} D \subset \mathcal{A}(K, C) \subset \bigcap_{D \in \mathcal{I}_{\mathcal{A}}(K, C)} D \subset K \quad (1)$$

We thus derive the following obvious but important statement:

Proposition 2.3 *Let us consider a map $\mathcal{A} : \mathcal{D}^2 \mapsto \mathcal{D}$.*

1. *If the map $D \mapsto \mathcal{A}(D, C)$ is a pre-opening and if $\mathcal{A}(K, C)$ belongs to the family $\mathcal{K}_{\mathcal{A}}(K, C)$, then $\mathcal{A}(K, C)$ is the largest fixed-point of the map $D \mapsto \mathcal{A}(D, C)$ between C and K . Furthermore, the map $D \mapsto \mathcal{A}(D, C)$ is an opening.*
2. *If the map $D \mapsto \mathcal{A}(K, D)$ is a pre-closing and if $\mathcal{A}(K, C)$ belongs to the family $\mathcal{I}_{\mathcal{A}}(K, C)$, then $\mathcal{A}(K, C)$ is the smallest fixed-point of the map $D \mapsto \mathcal{A}(K, D)$ between C and K . Furthermore, the map $D \mapsto \mathcal{A}(D, C)$ is a closing.*
3. *If the map $(K, C) \mapsto \mathcal{A}(K, C)$ is a pre-opening with respect to K and a pre-closing with respect to C , then any minimax $D \in \mathcal{D}(K, C)$ of the map \mathcal{A} between C and K in the sense that:*

$$\mathcal{A}(D, C) = D = \mathcal{A}(K, D)$$

is unique and is equal to $\mathcal{A}(K, C)$.

Consequently, if $\mathcal{A}(K, C)$ belongs to the intersection $\mathcal{K}_{\mathcal{A}}(K, C) \cap \mathcal{I}_{\mathcal{A}}(K, C)$, then $\mathcal{A}(K, C)$ is the unique minimax of the map \mathcal{A} between C and K :

$$\mathcal{A}(\mathcal{A}(K, C), C) = \mathcal{A}(K, C) = \mathcal{A}(K, \mathcal{A}(K, C))$$

In this case, \mathcal{A} is an opening with respect to K and a closing with respect to C .

Proof — The two first statements are obvious consequence of Lemma 2.2.

Therefore any minimax $D \in \mathcal{K}_{\mathcal{A}}(K, C) \cap \mathcal{I}_{\mathcal{A}}(K, C)$ is necessarily equal to $\mathcal{A}(K, C)$ of K and C , because since D belongs to $\mathcal{I}_{\mathcal{A}}(K, C)$, $\mathcal{A}(K, C) \subset \mathcal{A}(K, D) \subset D$ and since D belongs to $\mathcal{K}_{\mathcal{A}}(K, C)$, then $D \subset \mathcal{A}(D, C) \subset \mathcal{A}(K, C)$. Hence $\mathcal{A}(K, C) = D$. \square

The above Proposition 7.1 provides the uniqueness of the minimax of the map \mathcal{A} that is a pre-opening with respect to K and a pre-closing with respect to C . The existence of this minimax must be checked in each specific example by proving the two following inclusions

$$\begin{cases} i) & \mathcal{A}(K, C) \subset \mathcal{A}(\mathcal{A}(K, C), C) \\ ii) & \mathcal{A}(\mathcal{A}(K, C), C) \subset \mathcal{A}(K, C) \end{cases}$$

2.2 Galois Transform

Let us assume that $\emptyset \in \mathcal{D}$ and that the map $C \mapsto \mathcal{A}(K, C)$ is a dilation in the sense that for any family I of indexes, we have

$$\mathcal{A}\left(K, \bigcup_{i \in I} C_i\right) = \bigcup_{i \in I} \mathcal{A}(K, C_i)$$

Therefore, for any $C \in \mathcal{D}(K, \emptyset)$, the family

$$D \in \mathcal{D}(K, \emptyset) \quad \text{such that} \quad \mathcal{A}(K, D) \subset C$$

is max-closed (stable under the union) and has a largest element because, \mathcal{A} being a dilation with respect to C ,

$$\mathcal{A}\left(K, \bigcup_{\mathcal{A}(K, D) \subset C} D\right) = \bigcup_{\mathcal{A}(K, D) \subset C} \mathcal{A}(K, D) \subset C$$

In the same way, if $X \in \mathcal{D}$ and if the map $K \mapsto \mathcal{A}(K, C)$ is an erosion in the sense that

$$\mathcal{A}\left(\bigcap_{i \in I} K_i, C\right) = \bigcap_{i \in I} \mathcal{A}(K_i, C)$$

then for any $K \in \mathcal{D}(X, C)$, the family

$$D \in \mathcal{D}(X, C) \quad \text{such that} \quad K \subset \mathcal{A}(D, C)$$

is min-closed (stable under the intersection) and has a smallest element because, \mathcal{A} being an erosion with respect to K ,

$$K \subset \mathcal{A}\left(K, \bigcap_{K \subset \mathcal{A}(D, C)} D\right) = \bigcap_{K \subset \mathcal{A}(D, C)} \mathcal{A}(D, C)$$

Definition 2.4 Let us assume that \mathcal{A} is a dilation with respect to C and that \emptyset belongs to \mathcal{D} . Then we shall say that the map $C \mapsto \mathcal{A}^*(K, C)$ defined by

$$\forall K \in \mathcal{D}, \forall C \in \mathcal{D}(K, \emptyset), \mathcal{A}^*(K, C) := \bigcup_{\{D \in \mathcal{D}(K, \emptyset) \text{ such that } \mathcal{A}(K, D) \subset C\}} D$$

is called the upper Galois transform (or the upper residual) of \mathcal{A} with respect to C .

If \mathcal{A} is an erosion with respect to K and X belongs to \mathcal{D} , then we shall say that the map $\mathcal{A}_* : K \mapsto \mathcal{A}_*(K, C)$ defined by

$$\forall C \in \mathcal{D}, \forall K \in \mathcal{D}(X, C), \mathcal{A}_*(K, C) := \bigcap_{D \in \mathcal{D}(X, K) \text{ such that } K \subset \mathcal{A}(D, C)} D$$

is called the lower Galois transform (or the lower residual) of \mathcal{A} with respect to K .

Proposition 2.5 Let us assume that \mathcal{A} is a dilation and that \emptyset belongs to \mathcal{D} .

1. The map \mathcal{A}^* is increasing with respect to C and the inclusions

$$\mathcal{A}(K, \mathcal{A}^*(K, C)) \subset C \text{ \& } C \subset \mathcal{A}^*(K, \mathcal{A}(K, C))$$

hold true and actually, $C \mapsto \mathcal{A}^*(K, C)$ is the unique increasing map Φ satisfying

$$\mathcal{A}(K, \Phi(C)) \subset C \text{ \& } C \subset \Phi(\mathcal{A}(K, C))$$

2. If \mathcal{A} is increasing with respect to C , then \mathcal{A}^* is an erosion with respect to C , and its lower Galois transform with respect to C coincides with \mathcal{A} :

$$(\mathcal{A}^*)_*(K, C) = \mathcal{A}(K, C)$$

3. If \mathcal{A} is extensive with respect to C , then $\mathcal{A}^*(K, C) \subset C$,
4. If \mathcal{A} is increasing and idempotent with respect to C , then

$$\mathcal{A}^*(K, C) = \bigcup_{\{D \in \mathcal{I}_{\mathcal{A}}(K, \emptyset) \text{ such that } D \subset C\}} D$$

is the largest element $D \in \mathcal{I}_{\mathcal{A}}(C, \emptyset)$ contained in C and thus, $\mathcal{A}(K, \mathcal{A}^*(K, C)) = \mathcal{A}^*(K, C)$.

Let us assume that \mathcal{A} is an erosion and that X belongs to \mathcal{D} .

1. The map \mathcal{A}^* is increasing with respect to K and the inclusions

$$K \subset \mathcal{A}(\mathcal{A}_*(K, C), C) \ \& \ \mathcal{A}_*(\mathcal{A}(K, C), C) \subset K$$

hold true; Actually, $K \mapsto \mathcal{A}_*(K, C)$ is the unique increasing map Ψ satisfying

$$K \subset \mathcal{A}(K, \Psi(K)) \ \& \ \Psi(\mathcal{A}(K, C)) \subset K$$

2. If \mathcal{A} is increasing with respect to K , then \mathcal{A}^* is a dilation with respect to K , and its upper Galois transform with respect to K coincides with \mathcal{A} :

$$(\mathcal{A}_*)^*(K, C) = \mathcal{A}(K, C)$$

3. If \mathcal{A} is idempotent with respect to K , then

$$\mathcal{A}_*(K, C) = \bigcap_{D \in \mathcal{K}_{\mathcal{A}}(X, C) \text{ such that } K \subset D} D$$

is the smallest element $D \in \mathcal{K}_{\mathcal{A}}(X, K)$ containing K and thus, $\mathcal{A}(\mathcal{A}_*(K, C), C) = \mathcal{A}^*(K, C)$.

Proof — The map \mathcal{A}^* is increasing with respect to C because the family of subsets D such that $\mathcal{A}(K, D) \subset C_1$ is contained in the family of subsets D such that $\mathcal{A}(K, D) \subset C_2$ whenever $C_1 \subset C_2$. Inclusion $\mathcal{A}(K, \mathcal{A}^*(K, C)) \subset C$ holds true by construction and replacing C by $\mathcal{A}(K, C)$ in the definition of the upper Galois transform, we obtain the second inclusion $C \subset \mathcal{A}^*(K, \mathcal{A}(K, C))$.

Let Φ satisfy

$$\mathcal{A}(K, \Phi(C)) \subset C \ \& \ C \subset \Phi(\mathcal{A}(K, C))$$

Whenever $\mathcal{A}(K, D) \subset C$, then $D \subset \Phi(\mathcal{A}(K, D)) \subset \Phi(C)$ since Φ is increasing, and thus, $\mathcal{A}^*(K, C) \subset \Phi(C)$. Furthermore, inclusion $\mathcal{A}(K, \Phi(C)) \subset C$ implies that $\Phi(C) \subset \mathcal{A}^*(K, C)$.

Let us check now that \mathcal{A}^* is an erosion with respect to C . Indeed, since \mathcal{A}^* is increasing with respect to C , then $\mathcal{A}^*(K, \bigcap_{i \in I} C_i) \subset \bigcap_{i \in I} \mathcal{A}^*(K, C_i) =: D^*$. On the other hand, since for every $i \in I$, $D^* \subset \mathcal{A}^*(K, C_i) \subset C_i$, we infer that $D^* \subset \mathcal{A}^*(K, \bigcap_{i \in I} C_i)$.

Since \mathcal{A}^* is an erosion with respect to C , we can take its lower Galois transform defined by

$$(\mathcal{A}^*)_*(K, C) := \bigcap_{D \mid C \subset \mathcal{A}^*(K, D)} D$$

Since $C \subset \mathcal{A}^*(K, \mathcal{A}(K, C))$, we infer that $(\mathcal{A}^*)_*(K, C) \subset \mathcal{A}(K, C)$ by taking $D := \mathcal{A}(K, C)$. On the other hand, for any D satisfying $C \subset \mathcal{A}^*(K, D)$, we infer that $\mathcal{A}(K, C) \subset \mathcal{A}(K, \mathcal{A}^*(K, D))$

since \mathcal{A} is increasing with respect to C and we know that $\mathcal{A}(K, \mathcal{A}^*(K, D)) \subset D$. Hence $\mathcal{A}(K, C) \subset D$ for any D satisfying $C \subset \mathcal{A}^*(K, D)$, and thus, $\mathcal{A}(K, C) \subset \mathcal{A}^*(K, \mathcal{A}(K, C))$, so that $\mathcal{A}(K, C) = \mathcal{A}^*(K, \mathcal{A}(K, C))$.

Finally, assume that \mathcal{A} is increasing and idempotent with respect to C and prove that $\mathcal{A}(K, C)$ is the largest element $D \in \mathcal{I}_{\mathcal{A}}(K, \emptyset)$. First, we observe that inclusion

$$D^* := \bigcup_{D \in \mathcal{I}_{\mathcal{A}}(K, \emptyset) \mid D \subset C} D = \bigcup_{D \in \mathcal{I}_{\mathcal{A}}(K, \emptyset) \mid \mathcal{A}(K, D) \subset C} D \subset \mathcal{A}^*(K, C)$$

holds always true. On the other hand, if \mathcal{A} is idempotent with respect to C , we deduce from inclusions

$$\mathcal{A}(K, \mathcal{A}(K, \mathcal{A}^*(K, C))) = \mathcal{A}(K, \mathcal{A}^*(K, C)) \subset D$$

that

$$\mathcal{A}(K, \mathcal{A}^*(K, C)) \subset \mathcal{A}^*(K, C)$$

and thus, that $\mathcal{A}(K, \mathcal{A}^*(K, C)) \subset D^*$. \square

We deduce the following consequence:

Theorem 2.6 *If $\emptyset \in \mathcal{D}$ and if $C \mapsto \mathcal{A}(K, C)$ is both a closing and a dilation, then its upper Galois transform $C \mapsto \mathcal{A}^*(K, C)$ is both an opening and an erosion associating with any C the largest subset $D \in \mathcal{I}_{\mathcal{A}}(K, \emptyset)$ contained in C .*

Symmetrically, if $X \in \mathcal{D}$ and if $K \mapsto \mathcal{A}(K, C)$ is both an opening and an erosion, then its lower Galois transform $K \mapsto \mathcal{A}_(K, C)$ is both a closing and a dilation associating with any K the smallest subset $D \in \mathcal{K}_{\mathcal{A}}(X, C)$ containing K .*

Remark — We observe that

1. $\mathcal{A}(K, \emptyset)$ is the smallest subset $D \in \mathcal{I}_{\mathcal{A}}(K, \emptyset)$ contained in K ,
2. $\mathcal{A}^*(K, K)$ is the largest subset $D \in \mathcal{I}_{\mathcal{A}}(K, \emptyset)$ contained in K ,

so that, for any $C \subset K$,

$$\mathcal{A}(K, \emptyset) \subset \mathcal{A}(K, C) \subset \mathcal{A}^*(K, K) \subset K$$

In the same way,

1. $\mathcal{A}(X, C)$ is the largest subset $D \in \mathcal{K}_{\mathcal{A}}(X, C)$ containing K ,
2. $\mathcal{A}_*(C, C)$ is the smallest subset $D \in \mathcal{K}_{\mathcal{A}}(X, C)$ containing C ,

so that, for any $C \subset K$,

$$C \subset \mathcal{A}_*(C, C) \subset \mathcal{A}(K, C) \subset \mathcal{A}(X, C) \quad \square$$

3 Minimax Characterization of Viability Kernels and Capture Basins

We shall prove in this section the statement announced in the introduction that the viability kernel $\text{Viab}_F(K, C)$ of K with target C is the unique “minimax (bilateral fixed-point)” D between C and K of the map $(K, C) \mapsto \text{Viab}_F(K, C)$ in the sense that

$$C \subset D \subset K \ \& \ D = \text{Viab}(K, D) = \text{Viab}(D, C)$$

3.1 Evolutionary Systems

Actually, such results depend only on few properties of the solution map $x \rightsquigarrow \mathcal{S}_F(x) \subset \mathcal{C}(0, \infty; X)$ associating with any x the set $\mathcal{S}_F(x)$ of solutions to differential inclusions $x' \in F(x)$ starting at x , that are shared by solution maps of

1. control problems with memory (see the contributions of [47, 48, 49, Haddad] and [21, Aubin & Haddad], some of them being presented in [2, Aubin]) — before known under the name of functional control problems, the new fashion calling them as “path dependent control systems”
2. parabolic type partial differential inclusions (see the contributions of [65, 66, 67, Shi Shuzhong], some of them being presented in [2, Aubin]) — also known as distributed control systems
3. “mutational equations” governing the evolution in metric spaces, including “morphological equations” governing the evolution of sets (see [4, Aubin] for instance).

Such solution maps define evolutionary system:

Definition 3.1 *We regard $\mathcal{C}(0, \infty; X)$ as the space of evolutions. An evolutionary system is a set-valued map $\mathcal{S} : X \rightsquigarrow \mathcal{C}(0, \infty; X)$ satisfying*

1. *the initial property: $x(0) = x$,*
2. *the translation property: Let $x(\cdot) \in \mathcal{S}(x)$. Then for all $T \geq 0$, the function $y(\cdot)$ defined by $y(t) := x(t + T)$ is a solution $y(\cdot) \in \mathcal{S}(x(T))$ starting at $x(T)$,*
3. *the concatenation property: Let $x(\cdot) \in \mathcal{S}(x)$ and $T \geq 0$. Then for every $y(\cdot) \in \mathcal{S}(x(T))$, the function $z(\cdot)$ defined by*

$$z(t) := \begin{cases} x(t) & \text{if } t \in [0, T] \\ y(t - T) & \text{if } t \geq T \end{cases}$$

belongs to $\mathcal{S}(x)$.

3.2 Viability Kernels and Capture Basins

Definition 3.2 We say that an evolution $x(\cdot)$ is viable in K if for any $t \geq 0$, $x(t) \in K$ and we shall denote by $\mathcal{S}^K(x)$ the set of evolutions of the evolutionary system \mathcal{S} starting from $x \in K$ viable in K .

The viability kernel

$$\text{Viab}(K, C) := \text{Viab}_{\mathcal{S}}(K, C)$$

of K with target C under the evolutionary system \mathcal{S} is the subset of initial states $x \in K$ from which starts at least one evolution viable in K forever or until it reaches the target C in finite time.

The viable-capture basin

$$\text{Capt}(K, C) := \text{Capt}_{\mathcal{S}}(K, C)$$

of C viable in K under the evolutionary system \mathcal{S} is the subset of initial states $x \in K$ from which starts at least one evolution viable in K until it reaches the target C in finite time.

When the target $C = \emptyset$ is empty, we say that

$$\text{Viab}(K) := \text{Viab}_{\mathcal{S}}(K) := \text{Viab}_{\mathcal{S}}(K, \emptyset)$$

is the viability kernel of K under the evolutionary system.

We shall say that

1. a subset K is viable outside a target $C \subset K$ under the evolutionary system \mathcal{S} if $K := \text{Viab}(K, C)$, i.e., if from any $x \in K$ starts at least one evolution viable in K forever or until it reaches the target C in finite time,
2. a subset K captures the target $C \subset K$ in K under the evolutionary system \mathcal{S} if $K := \text{Capt}(K, C)$, i.e., from any $x \in K$ starts at least one evolution viable in K until it reaches the target C in finite time,
3. that C is isolated in K if $C = \text{Viab}(K, C)$,
4. that K is a repeller if $\text{Viab}(K) = \emptyset$, i.e., if the empty set is isolated in K .

Naturally,

$$C \subset \text{Capt}(K, C) \subset \text{Viab}(K, C) \subset K$$

3.3 Minimax Characterization of Viability Kernels

Let us consider a target $C \subset K$ and the family $\mathcal{D}(K, C)$ of subsets D satisfying $C \subset D \subset K$.

We associate with these subsets the two maps $\text{Viab}(K, \cdot) : \mathcal{D}(K, C) \mapsto \mathcal{D}(K, C)$ and $\text{Viab}(\cdot, C) : \mathcal{D}(K, C) \mapsto \mathcal{D}(K, C)$ associating with any subset $D \in \mathcal{D}(K, C)$ the viability kernels $\text{Viab}(K, D)$ and $\text{Viab}(D, C)$.

The family $\mathcal{K}_{\text{Viab}}(K, C)$ is then the family of subsets $D \in \mathcal{D}(K, C)$ viable outside C and the family $\mathcal{I}_{\text{Viab}}(K, C)$ is then the family of subsets $D \in \mathcal{D}(K, C)$ isolated in K .

Theorem 3.3 *The viability kernel $\text{Viab}(K, C)$ of a subset K with target $C \subset K$ is*

1. *the largest subset $D \in \mathcal{D}(K, C)$ viable outside the target C ,*
2. *the smallest subset $D \in \mathcal{D}(K, C)$ isolated in K ,*
3. *the unique minimax (bilateral fixed-point) $D \in \mathcal{D}(K, C)$ in the sense that*

$$D = \text{Viab}(K, D) = \text{Viab}(D, C)$$

It is antiextensive with respect to K and extensive with respect to C :

$$C \subset \text{Viab}(K, C) \subset K$$

increasing in the sense that

$$\text{If } C_1 \subset C_2 \text{ \& } K_1 \subset K_2, \text{ then } \text{Viab}(K_1, C_1) \subset \text{Viab}(K_2, C_2)$$

is an opening with respect to K , a closing with respect to C and the map $C \mapsto \text{Viab}(K, C)$ is a dilation in the sense that

$$\text{Viab} \left(K, \bigcup_{i \in I} C_i \right) = \bigcup_{i \in I} \text{Viab}(K, C_i)$$

The upper Galois transform $C \mapsto \text{Viab}^(K, C)$ of $C \mapsto \text{Viab}(K, C)$ is an opening and an erosion associating with $C \subset K$ the largest subset $\text{Viab}^*(K, C)$ contained in C and isolated in K and $\text{Viab}^*(K, K)$ is the largest subset of K isolated in K .*

The same properties hold true for the viable-capture map $\text{Capt}(K, C)$ of a target C viable in K .

Proof — The viability kernel map is obviously a pre-opening with respect to K and a pre-closing with respect to C . Therefore, thanks to Proposition 2.3, it is sufficient to prove that $\text{Viab}(K, C)$ belongs to the intersection $\mathcal{K}_{\text{Viab}}(K, C) \cap \mathcal{I}_{\text{Viab}}(K, C)$, i.e., that it is viable in outside C and isolated in K .

1. The translation property implies that the viability kernel $\text{Viab}(K, C)$ is viable outside C :

$$\text{Viab}(K, C) \subset \text{Viab}(\text{Viab}(K, C), C)$$

Take $x_0 \in \text{Viab}(K, C)$ and prove that there exists a solution $x(\cdot) \in \mathcal{S}(x_0)$ starting at x_0 viable in $\text{Viab}(K, C)$ until it possibly reaches C . Indeed, there exists a solution $x(\cdot) \in \mathcal{S}(x_0)$ viable in K until some time $T \geq 0$ either finite when it reaches C or infinite. Then for all $t \in [0, T[$, the function $y(\cdot)$ defined by $y(\tau) := x(t + \tau)$ is a solution $y(\cdot) \in \mathcal{S}(x(t))$ starting at $x(t)$ and viable in K until it reaches C at time $T - t$. Hence $x(t)$ does belong to $\text{Viab}(K, C)$ for every $t \in [0, T[$.

2. The concatenation property implies that the viability kernel $\text{Viab}(K, C)$ is isolated in K :

$$\text{Viab}(K, \text{Viab}(K, C)) \subset \text{Viab}(K, C)$$

Let x belongs to $\text{Viab}(K, \text{Viab}(K, C))$. There exists at least one evolution $x(\cdot) \in \mathcal{S}(x)$ would either remain in K or reach the viability kernel $\text{Viab}(K, C)$ in finite time. In this case, it can be concatenated with an evolution either remaining in $\text{Viab}(K, C) \subset K$ or reaching the target C in finite time. This implies that $x \in \text{Viab}(K, C)$.

The properties of the upper Galois transform $C \mapsto \text{Viab}^*(K, C)$ follow from Theorem 2.6.

The same proof shows that viable-capture basins always enjoy these properties. \square

Let us introduce the following notations. We associate with a subset D , a time $t \geq 0$ and a function $x(\cdot)$ the subset

$$J_D(t; x(\cdot)) := \begin{cases} x(0) & \text{if } x(t) \in D \\ \emptyset & \text{if } x(t) \notin D \end{cases}$$

and

$$I_D(t; x(\cdot)) := \bigcap_{s \in [0, t]} J(s, x(\cdot))$$

We next associate with a pair (K, C) of subset where $C \subset K$ the subset

$$L_{(K, C)}(t; x(\cdot)) := I_K(t; x(\cdot)) \cap J_C(t; x(\cdot))$$

Lemma 3.4 *The capture basin of C viable in K under the evolutionary system \mathcal{S} is equal to*

$$\text{Capt}(K, C) = \bigcup_{x \in K} \bigcup_{x(\cdot) \in \mathcal{S}(x)} \bigcup_{t \geq 0} L_{(K, C)}(t; x(\cdot)) \quad (2)$$

and thus, the unique fixed set of the system

$$\begin{cases} D = \bigcup_{x \in K} \bigcup_{x(\cdot) \in \mathcal{S}(x)} \bigcup_{t \geq 0} L_{(D,C)}(t; x(\cdot)) \\ = \bigcup_{x \in K} \bigcup_{x(\cdot) \in \mathcal{S}(x)} \bigcup_{t \geq 0} L_{(K,D)}(t; x(\cdot)) \end{cases} \quad (3)$$

Proof — To say that x belongs to the capture basin $\text{Capt}(K, C)$ of the target C viable in K amounts to saying that there exist at least one solution $x(\cdot) \in \mathcal{S}(x)$ and $t^* \geq 0$ such that

$$\begin{cases} i) & \forall s \in [0, t^*], \quad x(s) \in K \\ ii) & \quad \quad \quad \quad \quad x(t^*) \in C \end{cases}$$

i.e., if $x = x(0)$ belongs to $L_{(K,C)}(t^*, x(\cdot))$. In other words, x belongs to $\text{Capt}(K, C)$ if and only if

$$x \in \bigcup_{x(\cdot) \in \mathcal{S}(x)} \bigcup_{t \geq 0} L_{(K,C)}(t, x(\cdot))$$

Therefore formula (2) is proved. Using it, Theorem 3.3 can be reformulated as stating that the capture basin is the unique subset D satisfying (3). \square

4 Minimax Characterization of Invariance Kernels and Absorption Basins

4.1 Invariance Kernels and Absorption Basins

Definition 4.1 *The invariance kernel*

$$\text{Inv}(K, C) := \text{Inv}_{\mathcal{S}}(K, C)$$

of K with target C under the evolutionary system \mathcal{S} is the subset of initial states $x \in K$ such that all evolutions starting from x are viable in K forever or until they reach the target C in finite time.

The invariant-absorption basin

$$\text{Abs}(K, C) := \text{Abs}_{\mathcal{S}}(K, C)$$

of C invariant in K under the evolutionary system \mathcal{S} is the subset of initial states $x \in K$ such that all evolutions starting from x are viable in K until they reach the target C in finite time.

When the target $C = \emptyset$ is empty, we say that

$$\text{Inv}(K) := \text{Inv}_{\mathcal{S}}(K) := \text{Inv}_{\mathcal{S}}(K, \emptyset)$$

is the invariance kernel of K under the evolutionary system.

We shall say that

1. a subset K is invariant outside a target $C \subset K$ under the evolutionary system \mathcal{S} if $K := \text{Inv}(K, C)$, i.e., if from any $x \in K$ such that all evolutions starting from x are viable in K forever or until they reach the target C in finite time,
2. a subset K absorbs the target $C \subset K$ in K under the evolutionary system \mathcal{S} if $K := \text{Capt}(K, C)$, i.e., from any $x \in K$, such that all evolutions starting from x are viable in K until they reach the target C in finite time,
3. that C is separated in K if $C = \text{Inv}(K, C)$,

Naturally,

$$C \subset \text{Abs}(K, C) \subset \text{Inv}(K, C) \subset K$$

4.2 Minimax Characterization

We associate with these subsets the two maps $\text{Inv}(K, \cdot) : \mathcal{D}(K, C) \mapsto \mathcal{D}(K, C)$ and $\text{Inv}(\cdot, C) : \mathcal{D}(K, C) \mapsto \mathcal{D}(K, C)$ associating with any subset $D \in \mathcal{D}(K, C)$ the invariance kernels $\text{Inv}(K, D)$ and $\text{Inv}(D, C)$.

The family $\mathcal{K}_{\text{Inv}}(K, C)$ is then the family of subsets $D \in \mathcal{D}(K, C)$ invariant outside C and the family $\mathcal{I}_{\text{Inv}}(K, C)$ is then the family of subsets $D \in \mathcal{D}(K, C)$ separated in K .

Theorem 4.2 *The invariance kernel $\text{Inv}(K, C)$ of a subset K with target $C \subset K$ is*

1. the largest subset $D \in \mathcal{D}(K, C)$ invariant outside the target C ,
2. the smallest subset $D \in \mathcal{D}(K, C)$ separated in K ,
3. the unique minimax (bilateral fixed-point) $D \in \mathcal{D}(K, C)$ in the sense that

$$D = \text{Inv}(K, D) = \text{Inv}(D, C)$$

It is antiextensive with respect to K and extensive with respect to C :

$$C \subset \text{Inv}(K, C) \subset K$$

increasing in the sense that

$$\text{If } C_1 \subset C_2 \text{ \& } K_1 \subset K_2, \text{ then } \text{Inv}(K_1, C_1) \subset \text{Inv}(K_2, C_2)$$

is an opening with respect to K , a closing with respect to C and the map $C \mapsto \text{Inv}(K, C)$ is an erosion in the sense that

$$\text{Inv}\left(\bigcap_{i \in I} K_i, C\right) = \bigcap_{i \in I} \text{Inv}(K_i, C)$$

The lower Galois transform $K \mapsto \text{Inv}_*(K, C)$ of $K \mapsto \text{Inv}(K, C)$ is a closing and a dilation associating with K the smallest subset $\text{Inv}_*(K, C)$ containing K and invariant outside C , and $\text{Inv}_*(C, C)$ is the smallest invariant subset containing C . The same properties hold true for the invariant-absorption map $\text{Abs}(K, C)$ of a target C invariant in K .

We recognize the definition of invariance envelopes in the last statement of the above theorem:

Definition 4.3 We shall say that the smallest invariant subset containing C , equal to $\text{Inv}_*(C, C)$, is the invariance envelope of C and that the smallest subset of K containing C invariant outside C , equal to $\text{Inv}_*(K, C)$, is the invariance envelope of K outside C .

Proof — The invariance kernel map is obviously a pre-opening with respect to K and a pre-closing with respect to C . Therefore, thanks to Proposition 2.3, it is sufficient to prove that $\text{Inv}(K, C)$ belongs to the intersection $\mathcal{K}_{\text{Inv}}(K, C) \cap \mathcal{I}_{\text{Inv}}(K, C)$, i.e., that it is invariant outside C and separated in K .

1. The invariance kernel $\text{Inv}(K, C)$ is invariant outside C :

$$\text{Inv}(K, C) \subset \text{Inv}(\text{Inv}(K, C), C)$$

Take $x_0 \in \text{Inv}(K, C)$ and prove all solutions $x(\cdot) \in \mathcal{S}(x_0)$ starting at x_0 are viable in $\text{Inv}(K, C)$ until they possibly reach C . Indeed, take any solution $x(\cdot) \in \mathcal{S}(x_0)$, which is viable in K until some time $T \geq 0$ either finite when it reaches C or infinite. Then for all $t \in [0, T[$, consider any solution $y(\cdot) \in \mathcal{S}(x(t))$ starting at $x(t)$. Then the solution $z(\cdot)$ defined by

$$z(\tau) := \begin{cases} x(\tau) & \text{if } \tau \in [0, t] \\ y(\tau - t) & \text{if } \tau \geq t \end{cases}$$

is also a solution of \mathcal{S} starting at x_0 at time 0, and thus viable in K until it possibly reaches the target C at time $T - t$ since x_0 belongs to the invariance kernel $\text{Inv}(K, C)$ of K with target C . Therefore, for every $s \in [0, T]$, $y(s) = z(t + s) \in K$ and thus, $x(t)$ does belong to $\text{Inv}(K, C)$ for every $t \in [0, T[$.

2. The invariance kernel $\text{Inv}(K, C)$ is separated from K :

$$\text{Inv}(K, \text{Inv}(K, C)) \subset \text{Inv}(K, C)$$

Let x belongs to $\text{Inv}(K, \text{Inv}(K, C))$. All evolutions $x(\cdot) \in \mathcal{S}(x)$ would either remain in K or reach the invariance kernel $\text{Inv}(K, C)$ in finite time T at some $x(T) \in \text{Inv}(K, C)$. Therefore, the translation $y(\tau) := x(\tau + T)$ is an evolution starting from $x(T)$, and thus, either remains in $\text{Inv}(K, C) \subset K$ or reaches the target C in finite time $S \geq 0$. This implies that for all $t \geq T$, $x(t) = y(t - T)$ is viable in K for ever or reach the target C at time $T + S$, i.e., that x_0 belongs to $\text{Inv}(K, C)$.

The properties of the lower Galois transform $K \mapsto \text{Inv}_*(K, C)$ follow from Theorem 2.6.

The same proof shows that invariant-absorption basins also enjoy these properties. \square

Lemma 4.4 *The Absorption basin of C invariant in K under the evolutionary system \mathcal{S} is equal to*

$$\text{Abs}(K, C) = \bigcup_{x \in K} \bigcap_{x(\cdot) \in \mathcal{S}(x)} \bigcup_{t \geq 0} L_{(K, C)}(t; x(\cdot)) \quad (4)$$

and thus, the unique fixed set of the system

$$\begin{cases} D = \bigcup_{x \in K} \bigcap_{x(\cdot) \in \mathcal{S}(x)} \bigcup_{t \geq 0} L_{(D, C)}(t; x(\cdot)) \\ = \bigcup_{x \in K} \bigcap_{x(\cdot) \in \mathcal{S}(x)} \bigcup_{t \geq 0} L_{(K, D)}(t; x(\cdot)) \end{cases} \quad (5)$$

Proof — By definition, x belongs to the absorption basin $\text{Abs}(K, C)$ of the target C invariant in K , for all solutions $x(\cdot) \in \mathcal{S}(x)$, there exist $t^* \geq 0$ such that

$$\begin{cases} i) & \forall s \in [0, t^*], \quad x(s) \in K \\ ii) & \quad \quad \quad \quad \quad x(t^*) \in C \end{cases}$$

i.e., if $x = x(0)$ belongs to $L_{(K, C)}(t^*, x(\cdot))$. In other words, x belongs to $\text{Abs}(K, C)$ if and only if

$$x \in \bigcap_{x(\cdot) \in \mathcal{S}(x)} \bigcup_{t \geq 0} L_{(K, C)}(t, x(\cdot))$$

Therefore formula (4) is proved. Using it, Theorem 4.2 can be reformulated as stating that the absorption basin is the unique subset D satisfying (5). \square

5 Construction of Openings and Closings

When the map \mathcal{A} is a pre-opening with respect to K , but not necessarily an opening, or/and a pre-closing with respect to C , but not necessarily a closing, Lemma 2.2 suggests to associate with the map \mathcal{A} the maps \mathcal{A}^\sharp and \mathcal{A}^\flat defined by

$$\mathcal{A}^\sharp(K, C) := \bigcup_{D \in \mathcal{K}_{\mathcal{A}}(K, C)} D \quad \& \quad \mathcal{A}^\flat(K, C) := \bigcap_{D \in \mathcal{I}_{\mathcal{A}}(K, C)} D$$

5.1 Definition of Associated Openings and Closings

We observe that if \mathcal{A} is a pre-opening with respect to K , then

$$C \subset \mathcal{A}^\sharp(K, C) \subset \mathcal{A}(K, C) \subset K$$

and thus, that

$$\forall D \in \mathcal{K}_{\mathcal{A}}(K, C), \quad \mathcal{A}(D, C) = \mathcal{A}^\flat(D, C) = D$$

Symmetrically, if \mathcal{A} is a pre-closing with respect to C , then

$$C \subset \mathcal{A}(K, C) \subset \mathcal{A}^\flat(K, C) \subset K$$

and

$$\forall D \in \mathcal{I}_{\mathcal{A}}(K, C), \quad \mathcal{A}(D, C) = \mathcal{A}^\sharp(D, C) = D$$

Consequently, if the map $(K, C) \mapsto \mathcal{A}(K, C)$ is a pre-opening with respect to K and a pre-closing with respect to C , then inequalities (1) can be written in the form

$$\forall (K, C) \in \mathcal{D}^2, \quad C \subset \mathcal{A}^\sharp(K, C) \subset \mathcal{A}(K, C) \subset \mathcal{A}^\flat(K, C) \subset K \quad (6)$$

Therefore, Lemma 2.2 states that *if \mathcal{A} is a pre-opening with respect to K and if $\mathcal{A}(K, C)$ belongs to $\mathcal{K}_{\mathcal{A}}(K, C)$, then $\mathcal{A}(K, C) = \mathcal{A}^\sharp(K, C)$ and that if \mathcal{A} is a pre-closing with respect to C and if $\mathcal{A}(K, C)$ belongs to $\mathcal{I}_{\mathcal{A}}(K, C)$, then $\mathcal{A}(K, C) = \mathcal{A}^\flat(K, C)$.*

5.2 The Matheron Theorem

We begin by observing that the supremum $\mathcal{A}^\sharp(K, C)$ belongs to $\mathcal{K}_{\mathcal{A}}(K, C)$ and is actually the “maximum” and that $\mathcal{A}^\flat(K, C)$ belongs to $\mathcal{I}_{\mathcal{A}}(K, C)$ and is the minimum:

Proposition 5.1 *Assume that the family \mathcal{D} is max-closed and that the map $D \mapsto \mathcal{A}(D, C)$ is a pre-opening. Then the family $\mathcal{K}_{\mathcal{A}}(K, C)$ is max-closed (stable for the union) and $\mathcal{A}^\sharp(K, C) \in \mathcal{K}_{\mathcal{A}}(K, C)$ is the largest fixed-point of the map $D \mapsto \mathcal{A}(D, C)$ in $\mathcal{D}(K, C)$:*

$$\mathcal{A}(\mathcal{A}^\sharp(K, C), C) = \mathcal{A}^\sharp(K, C)$$

Symmetrically, if the family \mathcal{D} is min-closed (stable for the intersection) and if the map $D \mapsto \mathcal{A}(K, D)$ is a pre-closing, then the family $\mathcal{I}_{\mathcal{A}}(K, C)$ is min-closed and $\mathcal{A}^b(K, C) \in \mathcal{I}_{\mathcal{A}}(K, C)$ is the smallest fixed-point of the map $D \mapsto \mathcal{A}(K, D)$ in $\mathcal{D}(K, C)$:

$$\mathcal{A}(K, \mathcal{A}^b(K, C)) = \mathcal{A}^b(K, C)$$

Proof — We reproduce the proof that can be found for instance in [55, Mattioli & Schmitt], Theorem III.3.3, p.46. Consider a family of subsets $D_i \in \mathcal{K}_{\mathcal{A}}(K, C)$. Therefore $D_i \subset \mathcal{A}(D_i, C) \subset \mathcal{A}(\bigcup_{i \in I} D_i, C)$ since \mathcal{A} is antiextensive and increasing. Consequently,

$$\bigcup_{i \in I} D_i \subset \mathcal{A}\left(\bigcup_{i \in I} D_i, C\right)$$

so that $\bigcup_{i \in I} D_i$ belongs to $\mathcal{K}_{\mathcal{A}}(K, C)$. This proves that the family $\mathcal{K}_{\mathcal{A}}(K, C)$ is stable for the union. Hence its supremum $\mathcal{A}^\#(K, C)$ belongs to $\mathcal{K}_{\mathcal{A}}(K, C)$ and thus, satisfies

$$\mathcal{A}(\mathcal{A}^\#(K, C), C) = \mathcal{A}^\#(K, C)$$

If $\mathcal{A}(K, C)$ belongs to $\mathcal{K}_{\mathcal{A}}(K, C)$, then

$$\mathcal{A}(K, C) \subset \mathcal{A}(\mathcal{A}(K, C), C) \subset \mathcal{A}^\#(K, C) \subset \mathcal{A}(K, C) \quad \square$$

Matheron's theorem (see [52, Matheron']) states in essence that the map $D \mapsto \mathcal{A}^\#(D, C)$ is an opening and that the map $D \mapsto \mathcal{A}^b(K, D)$ is a closing:

Theorem 5.2 (Matheron) *Assume that the family \mathcal{D} is max-closed and that the map $D \mapsto \mathcal{A}(D, C)$ is a pre-opening and $D \mapsto \mathcal{A}^\#(K, D)$ is extensive whenever $D \mapsto \mathcal{A}(K, D)$ is extensive¹. Then the map $D \mapsto \mathcal{A}^\#(D, C)$ is the unique opening onto the family $\mathcal{K}_{\mathcal{A}}(K, C)$:*

$$\mathcal{A}^\#(\mathcal{A}^\#(K, C), C) = \mathcal{A}^\#(K, C)$$

and the families

$$\mathcal{K}_{\mathcal{A}}(K, C) = \mathcal{K}_{\mathcal{A}^\#}(K, C)$$

coincide.

Similarly, if the family \mathcal{D} is min-closed and if the map $D \mapsto \mathcal{A}(K, D)$ is a pre-closing, then the map $D \mapsto \mathcal{A}^b(K, D)$ is the unique closing onto the family $\mathcal{I}_{\mathcal{A}}(K, C)$:

$$\mathcal{A}^b(K, \mathcal{A}^b(K, C)) = \mathcal{A}^b(K, C)$$

and the families

$$\mathcal{I}_{\mathcal{A}}(K, C) = \mathcal{I}_{\mathcal{A}^b}(K, C)$$

coincide.

¹However, $D \mapsto \mathcal{A}^\#(K, D)$ is not necessarily increasing whenever $D \mapsto \mathcal{A}(K, D)$ is increasing, although this happens for some specific examples.

Proof — We reproduce the proof that can be found for instance in [52, Matheron] and [55, Mattioli & Schmitt], Theorem III.3.3, p.46. The map $K \mapsto \mathcal{A}^\sharp(K, C) := \bigcup_{D \in \mathcal{K}_{\mathcal{A}}(K, C)} D \subset K$ is antiextensive. Since $\mathcal{K}_{\mathcal{A}}(K_1, C) \subset \mathcal{K}_{\mathcal{A}}(K_2, C)$ whenever $K_1 \subset K_2$, the map $K \mapsto \mathcal{A}^\sharp(K, C)$ is increasing.

If D belongs to $\mathcal{K}_{\mathcal{A}}(K, C)$, then $D \subset \mathcal{A}^\sharp(D, C)$ and thus belongs to $\mathcal{K}_{\mathcal{A}^\sharp}(K, C)$. Taking $D := \mathcal{A}^\sharp(K, C)$ that belongs to $\mathcal{K}_{\mathcal{A}^\sharp}(K, C)$, we infer that

$$\mathcal{A}^\sharp(K, C) \subset \mathcal{A}^\sharp(\mathcal{A}^\sharp(K, C), C)$$

By Proposition 2.3, $\mathcal{A}^\sharp(K, C)$ is the largest fixed point $D \in \mathcal{K}_{\mathcal{A}^\sharp}(K, C)$ of the map $K \mapsto \mathcal{A}^\sharp(K, C)$. Hence this map is an opening.

Conversely, if D belongs to $\mathcal{K}_{\mathcal{A}^\sharp}(K, C)$, then

$$D \subset \mathcal{A}^\sharp(D, C) := \mathcal{A}(D, C)$$

so that $D \subset \mathcal{A}(D, C)$ and thus belongs to $\mathcal{K}_{\mathcal{A}}(K, C)$. Consequently, the families \mathcal{A} and \mathcal{A}^\sharp coincide:

$$\mathcal{K}_{\mathcal{A}}(K, C) = \mathcal{K}_{\mathcal{A}^\sharp}(K, C)$$

Let us consider now a surjective opening $\Pi : \mathcal{D}(K, C) \mapsto \mathcal{K}_{\mathcal{A}}(K, C)$. Since $\Pi(K)$ belongs to $\mathcal{K}_{\mathcal{A}}(K, C) = \mathcal{K}_{\mathcal{A}^\sharp}(K, C)$, then $\Pi(K) \subset \mathcal{A}^\sharp(\Pi(K), C)$. Since $\mathcal{A}^\sharp(\cdot, C)$ is increasing and Π is antiextensive, we deduce that $\mathcal{A}^\sharp(\Pi(K), C) \subset \mathcal{A}^\sharp(K, C)$. We finally check that $\mathcal{A}^\sharp(K, C) \subset \Pi(K)$ because, for every $D \in \mathcal{K}_{\mathcal{A}^\sharp}(K, C)$,

$$D \subset \mathcal{A}^\sharp(D, C) = \mathcal{A}^\sharp(\Pi(D), C) \subset \Pi(D) \subset \Pi(K)$$

since $\Pi(D) = D \subset \Pi(K)$ whenever $D \in \mathcal{K}_{\mathcal{A}}(K, C)$ and $\Pi(D) \subset \Pi(K)$. Therefore $\Pi(K) = \mathcal{A}^\sharp(K, C)$. \square

5.3 The Opening and Closing Algorithms

One can define the Opening and Closing Algorithms for computing the opening \mathcal{A}^\sharp and closing \mathcal{A}^\flat associated with \mathcal{A} .

For the closing \mathcal{A}^\flat , we define recursively the Closing Algorithm by a sequence of subsets C_n by $C_0 := C$ and

$$\forall n \geq 0, C_{n+1} := \mathcal{A}(K, C_n)$$

Then, if \mathcal{A} is increasing with respect to K ,

$$\bigcup_{n \geq 0} C_n \subset \mathcal{A}^\flat(K, C)$$

Indeed, let any D belong to $\mathcal{I}_{\mathcal{A}}(K, C)$. Hence $C_0 := C \subset D$, and thus $\mathcal{A}(K, D) \subset D$ and $C_1 := \mathcal{A}(K, C_0) \subset \mathcal{A}(K, D)$ since \mathcal{A} is increasing with respect to C , so that $C_1 \subset D$. Assume now that $C_n \subset D$ and check that $C_{n+1} \subset D$. We infer that $\mathcal{A}(K, D) \subset D$ and $C_{n+1} := \mathcal{A}(K, C_n) \subset \mathcal{A}(K, D)$, so that

$$\forall D \in \mathcal{I}_{\mathcal{A}}(K, C), \bigcup_{n \geq 0} C_n \subset D$$

In the same way, the Opening Algorithm for computing the closing \mathcal{A}^\sharp associated with \mathcal{A} is defined recursively by $K_0 := K$ and

$$\forall n \geq 0, K_{n+1} := \mathcal{A}(K_n, C)$$

Then, if \mathcal{A} is increasing with respect to K ,

$$\mathcal{A}^\sharp(K, C) \subset \bigcap_{n \geq 0} K_n$$

Indeed, let any D belong to $\mathcal{K}_{\mathcal{A}}(K, C)$. Hence $D \subset K_0 := K$, and thus $D \subset \mathcal{A}(D, C)$ and $\mathcal{A}(D, C) \subset \mathcal{A}(K_0, C) =: K_1$ since \mathcal{A} is increasing with respect to K . Assume now that $D \subset K_n$ and check that $D \subset K_{n+1}$. We infer that $D \subset \mathcal{A}(D, C)$ and $\mathcal{A}(D, C) \subset \mathcal{A}(K_n, C) =: K_{n+1}$, so that

$$\forall D \in \mathcal{K}_{\mathcal{A}}(K, C), D \subset \bigcap_{n \geq 0} K_n$$

6 Viability Kernels and Capture Basins Algorithms for Discrete Systems

Let us consider a set-valued map $\Phi : X \rightsquigarrow X$ from a metric space X to itself, governing the evolution $\vec{x} : n \mapsto x_n$ defined by

$$\forall n \geq 0, x_{n+1} \in \Phi(x_n)$$

We denote by $\mathcal{S}_\Phi(x)$ the set of evolutions \vec{x} of solutions to the above discrete system starting at x_0 . Replacing the space $\mathcal{C}(0, \infty; X)$ of continuous time-dependent functions by the space $X^{\mathbf{N}}$ of discrete-time dependent function (sequences) and making the necessary adjustments in definitions, we can still regard \mathcal{S}_Φ as an evolutionary system from X to $X^{\mathbf{N}}$. We thus define the viability kernels $\text{Viab}_{\mathcal{S}_\Phi}(K, C)$ and the invariance kernels $\text{Inv}_{\mathcal{S}_\Phi}(K, C)$ and derive from Theorems 3.3 and 4.2 the following

Theorem 6.1 *The viability kernel $\text{Viab}_{\mathcal{S}_\Phi}(K, C)$ of a subset K with target $C \subset K$ is the unique minimax (bilateral fixed-point) $D \in \mathcal{D}(K, C)$ in the sense that*

$$D = \text{Viab}_{\mathcal{S}(\oplus)}(K, D) = \text{Viab}_{\mathcal{S}(\oplus)}(D, C)$$

and the invariance kernel $\text{Inv}_{\mathcal{S}(\oplus)}(K, C)$ of a subset K with target $C \subset K$ is the unique minimax (bilateral fixed-point) $D \in \mathcal{D}(K, C)$ in the sense that

$$D = \text{Inv}_{\mathcal{S}(\oplus)}(K, D) = \text{Inv}_{\mathcal{S}(\oplus)}(D, C)$$

6.1 An Alternate Characterization of Viability and Invariance Kernels

We shall prove that the viability and invariance kernels under a discrete systems are fixed points of openings associated with maps \mathcal{A} that we shall define.

We set $\Phi^{-1}(K) := \{x \mid \Phi(x) \cap K \neq \emptyset\}$ and $\Phi^{\ominus 1}(K) := \{x \mid \Phi(x) \subset K\}$ and we introduce the maps $\mathcal{A}_{\Phi^{-1}} : \mathcal{D} \times \mathcal{D} \mapsto \mathcal{D}$ and $\mathcal{A}_{\Phi^{\ominus 1}}$ defined on the family \mathcal{D} of all subsets of X by

$$\begin{cases} i) & \mathcal{A}_{\Phi^{-1}}(K, C) := C \cup (K \cap \Phi^{-1}(K)) \\ ii) & \mathcal{A}_{\Phi^{\ominus 1}}(K, C) := C \cup (K \cap \Phi^{\ominus 1}(K)) \end{cases}$$

They are pre-openings with respect to K and a pre-closings with respect to C .

We then can go one step farther and characterize these maps as openings and closings of these associated maps : For that purpose, we provide a characterization of the discrete viability and invariance kernels:

Theorem 6.2 *The viability kernel $\text{Viab}_{\mathcal{S}_\Phi}(K, C)$ of K with target C under Φ is the largest subset D of the family $\mathcal{K}_{\mathcal{A}_{\Phi^{-1}}}(K, C)$ of subsets $D \in \mathcal{D}(K, C)$ such that $D \subset C \cup (D \cap \Phi^{-1}(D))$.*

In the same way, the invariance kernel $\text{Inv}_{\mathcal{S}_\Phi}(K, C)$ of K with target C under Φ is the largest subset D of the family $\mathcal{K}_{\mathcal{A}_{\Phi^{\ominus 1}}}(K, C)$ of subsets $D \in \mathcal{D}(K, C)$ such that $D \subset C \cup (D \cap \Phi^{\ominus 1}(D))$.

Proof — Indeed, to say that $D \subset C \cup (D \cap \Phi^{-1}(D))$ means that $D \setminus C \subset \Phi^{-1}(D)$, i.e., that for any $x \in D$, either x belongs to C or else, that there exists $y \in \Phi(x) \cap D$. Consequently, for any $x_0 \in D$, either $x_0 \in C$ and x_0 belongs to the viability kernel $\text{Viab}_{\mathcal{S}_\Phi}(K, C)$ of K with target C or else, there exists $x_1 \in \Phi(x_0) \cap D$, so that we can continue by induction constructing a solution $\vec{x} \in \mathcal{S}_\Phi(x_0)$ starting from $x_0 \in D$ viable in D — and thus, in K — either until some finite N when $x_N \in C$ or forever. Hence $D \subset \text{Viab}_{\mathcal{S}_\Phi}(D, C) \subset \text{Viab}_{\mathcal{S}_\Phi}(K, C)$.

Conversely, if $D \subset \text{Viab}_{\mathcal{S}_\Phi}(D, C)$, there exists $\vec{x} \in \mathcal{S}_\Phi(x_0)$ starting from x_0 and viable in D outside C . Therefore, starting from any $x_0 \in D$, either $x_0 \in C$ or else, there exists $x_1 \in \Phi(x_0)$ that belongs to D , i.e., $x_0 \in \Phi^{-1}(x_1) \subset \Phi^{-1}(D)$. Hence $D \subset C \cup (D \cap \Phi^{-1}(D))$. \square

The above Characterization Theorem 6.2 and Matheron's Theorem 5.2 imply the following result:

Theorem 6.3 *Let $\Phi : X \mapsto X$ be a set-valued map from a set X to itself.*

1. *The map $K \mapsto \text{Viab}_{\mathcal{S}_\Phi}(K, C)$ is the opening associated with the map $\mathcal{A}_{\Phi^{-1}} : K \mapsto C \cup (K \cap \Phi^{-1}(K))$,*
2. *The map $K \mapsto \text{Inv}_{\mathcal{S}_\Phi}(K, C)$ is the opening associated with the map $\mathcal{A}_{\Phi^{\ominus 1}} : K \mapsto C \cup (K \cap \Phi^{\ominus 1}(K))$.*

Proof — Indeed, Theorem 6.2 states that $\text{Viab}_{\mathcal{S}_\Phi}(K, C)$ is the largest of subsets $D \in \mathcal{K}_{\mathcal{A}_{\Phi^{-1}}}(K, C)$, and thus, Theorem 5.2 implies that

$$\text{Viab}_{\mathcal{S}_\Phi}(K, C) := \mathcal{A}_{\Phi^{-1}}^\#(K, C)$$

The proof for the invariance kernel is identical. \square

6.2 The Viability Kernel Algorithm

Since the discrete viability kernel map $K \mapsto \text{Viab}_{\mathcal{S}_\Phi}(K, C)$ is the opening associated with the pre-opening $K \mapsto C \cup (K \cap \Phi^{-1}(K))$, we can define the Opening Algorithm which, here, is the Viability Algorithm designed by Patrick Saint-Pierre in [61, Saint-Pierre] :

Starting with $K_0 := K$, we define recursively the subsets K_n by

$$\forall n \geq 0, K_{n+1} := C \cup (K_n \cap \Phi^{-1}(K_n))$$

We already know that

$$\text{Viab}_{\mathcal{S}_\Phi}(K, C) \subset \bigcap_{n \geq 0} K_n$$

Adequate topological assumptions imply the equality:

Proposition 6.4 *Let us assume that K and $C \subset K$ are closed and that the graphical restriction² $\Phi|_K^K$ of Φ to $K \times K$ is upper semicontact³. Then the subsets K_n are closed and*

$$\text{Viab}_{\mathcal{S}_\Phi}(K, C) = \bigcap_{n \geq 0} K_n$$

²defined by $\text{Graph}(\Phi|_K^K) := \text{Graph}(\Phi)$.

³A set-valued map $\Phi : X \rightsquigarrow Y$ is said to be *upper semicontact* at x if for every sequence $x_n \in \text{Dom}(\Phi)$ converging to x and for every sequence $y_n \in \Phi(x_n)$, there exists a subsequence y_{n_p} converging to some $y \in \Phi(x)$.

Proof — Let x belongs to $K_\infty := \bigcap_{n \geq 0} K_n$. If $x \in K_\infty \setminus C$, then for every $n \geq 0$, $x \in \Phi^{-1}(K_n)$, and thus, there exists $y_n \in \Phi(x) \in K_n$. Since the images $\Phi(x) \cap K$ are compact, a subsequence (again denoted by) y_n converges to some $y_* \in \Phi(x) \cap \bigcap_{n \geq 0} \overline{K_n}$. Hence, whenever the subsets K_n are closed,

$$x \in K_\infty \cap \Phi^{-1}(y_*) \subset K_\infty \cap \Phi^{-1}(K_\infty)$$

and thus, K_∞ is viable outside C under Φ , and consequently, contained in $\text{Viab}_{\mathcal{S}_\Phi}(K, C)$.

To prove recursively that the subsets K_n are closed, it is enough to prove that $K_{n+1} := K_n \cap \Phi^{-1}(K_n)$ is closed whenever K_n is closed. Let $x_j \in K_{n+1}$ converge to x , that belongs to K_n . Furthermore, there exists $y_j \in K_n \cap \Phi(x_j)$. Since the graphical restriction $\Phi|_K^K$ of Φ is assumed to be upper semicontact, a subsequence (again denoted by) y_j converges to $y \in \Phi|_K^K(x)$ and actually, to $\Phi(x) \in K_n$. This implies that K_{n+1} is closed. \square

For the invariance kernels, starting with $L_0 := K$, we define recursively the subsets L_n by

$$\forall n \geq 0, L_{n+1} := C \cup (L_n \cap \Phi^{\ominus 1}(L_n))$$

We already know that

$$\text{Inv}_{\mathcal{S}_\Phi}(K, C) \subset \bigcap_{n \geq 0} L_n$$

Proposition 6.5 *If we assume that K and $C \subset K$ are closed and that Φ is lower semicontinuous, then the subsets L_n are closed and*

$$\text{Inv}_{\mathcal{S}_\Phi}(K, C) = \bigcap_{n \geq 0} L_n$$

Proof — Let x belongs to $L_\infty := \bigcap_{n \geq 0} L_n$. If $x \in L_\infty \setminus C$, then for every $n \geq 0$, $x \in \Phi^{\ominus 1}(K_n)$, and thus, $\Phi(x_n) \subset K_n$. Hence

$$\Phi(x) \subset \bigcap_{n \geq 0} L_n =: L_\infty$$

Therefore,

$$x \in L_\infty \cap \Phi^{\ominus 1}(L_\infty)$$

and thus, L_∞ is invariant outside C under Φ , and consequently, contained in $\text{Inv}_{\mathcal{S}_\Phi}(K, C)$.

To prove recursively that the subsets L_n are closed, it is enough to prove that $L_{n+1} := L_n \cap \Phi^{\ominus 1}(L_n)$ is closed whenever L_n is closed. Let $x_j \in L_{n+1}$ converge to x , that belongs to

L_n . Since Φ is lower semicontinuous, for any $y \in \Phi(x)$, there exists $y_j \in \Phi(x_j)$ converging to y . Since $\Phi(x_j) \subset L_n$, we know that y_j belongs to L , and consequently, that y belongs to L_n . Hence $\Phi(x) \subset L_n$, so that x belongs to L_{n+1} . \square

6.3 An Alternate Characterization of Capture and Absorptions Basins

Let us consider now the capture basins $\text{Capt}_{\mathcal{S}_\Phi}(K, C)$ and the absorption basins $\text{Abs}_{\mathcal{S}_\Phi}(K, C)$ and provide another characterization involving the maps $\mathcal{B}_{\Phi^{-1}} : \mathcal{D} \times \mathcal{D} \mapsto \mathcal{D}$ and $\mathcal{B}_{\Phi^{\ominus 1}}$ defined by

$$\begin{cases} i) & \mathcal{B}_{\Phi^{-1}}(K, C) := K \cap (C \cup \Phi^{-1}(C)) \\ ii) & \mathcal{B}_{\Phi^{\ominus 1}}(K, C) := K \cap (C \cup \Phi^{\ominus 1}(C)) \end{cases}$$

They are pre-openings with respect to D and a pre-closings with respect to C .

Theorem 6.6 *The capture basin $\text{Capt}_{\mathcal{S}_\Phi}(K, C)$ of K of C viable in K under Φ is the smallest subset D of the family $\mathcal{I}_{\mathcal{B}_{\Phi^{-1}}}(K, C)$ of subsets $D \in \mathcal{D}(K, C)$ such that $K \cap (D \cup \Phi^{-1}(D)) \subset D$.*

In the same way, the absorption basin $\text{Abs}_{\mathcal{S}_\Phi}(K, C)$ of K with target C under Φ is the smallest subset D of the family $\mathcal{I}_{\mathcal{B}_{\Phi^{\ominus 1}}}(K, C)$ of subsets $D \in \mathcal{D}(K, C)$ such that $K \cap (D \cup \Phi^{\ominus 1}(D)) \subset D$.

Proof — The capture basin $\text{Capt}_{\mathcal{S}_\Phi}(K, C)$ belongs to the family $\mathcal{I}_{\mathcal{B}_{\Phi^{-1}}}(K, C)$. We have to check that whenever we take $x \in K \setminus \text{Capt}_{\mathcal{S}_\Phi}(K, C)$ satisfying

$$x \in \Phi^{-1}(\text{Capt}_{\mathcal{S}_\Phi}(K, C))$$

then such an x belongs to the capture basin. Indeed, such an element satisfies $\Phi(x) \cap \text{Capt}_{\mathcal{S}_\Phi}(K, C) \neq \emptyset$. Then x captures $\text{Capt}_{\mathcal{S}_\Phi}(K, C)$ in one step and thus, the target C in a finite number of steps. Hence

$$K \cap (\text{Capt}_{\mathcal{S}_\Phi}(K, C) \cup \Phi^{-1}(\text{Capt}_{\mathcal{S}_\Phi}(K, C))) \subset \text{Capt}_{\mathcal{S}_\Phi}(K, C)$$

Let us prove now that the capture basin is the smallest of the subsets $D \in \mathcal{I}_{\mathcal{B}_{\Phi^{-1}}}(K, C)$. Otherwise, there would exist some $D_0 \in \mathcal{I}_{\mathcal{B}_{\Phi^{-1}}}(K, C)$ and some

$$x_0 \in \text{Capt}_{\mathcal{S}_\Phi}(K, C) \setminus D_0$$

Taking the complement, we see that $K \setminus D_0 \subset \Phi^{\ominus 1}(X \setminus D_0)$. Since $x_0 \in K \setminus D_0$, we infer that $\Phi(x_0) \subset X \setminus D_0 \subset X \setminus C$, so that no solution $\vec{x} \in \mathcal{S}(x_0)$ can ever reach the target C before leaving K . \square

The above Characterization Theorem 6.2 and Matheron's Theorem 5.2 imply the following result:

Theorem 6.7 *Let $\Phi : X \mapsto X$ be a set-valued map from a set X to itself.*

1. *The map $C \mapsto \text{Capt}_{\mathcal{S}_\Phi}(K, C)$ is the closing associated with the map $C \mapsto K \cap (C \cup \Phi^{-1}(C))$,*
2. *The map $C \mapsto \text{Abs}_{\mathcal{S}_\Phi}(K, C)$ is the closing associated with the map $C \mapsto K \cap (C \cup \Phi^{\ominus 1}(C))$,*

Proof — Indeed, Theorem 6.6 states that $\text{Capt}_{\mathcal{S}_\Phi}(K, C)$ is the smallest of subsets $D \in \mathcal{I}_{\mathcal{B}_{\Phi^{-1}}}(K, C)$, and thus, Theorem 5.2 implies that

$$\text{Capt}_{\mathcal{S}_\Phi}(K, C) := \mathcal{B}_{\Phi^{-1}}^b(K, C)$$

The proof for the absorption basin is identical. \square

6.4 The Capture Basin Algorithm

Since the discrete capture basin map $C \mapsto \text{Capt}_{\mathcal{S}_\Phi}(K, C)$ is the closing associated with the pre-closing $C \mapsto K \cap (C \cup \Phi^{-1}(C))$, we can define the Closing Algorithm which, here, is the Capture Basin Algorithm designed by Patrick Saint-Pierre and Dominique Pujal :

Starting with $C_0 := C$, we define recursively the subsets C_n by

$$\forall n \geq 0, C_{n+1} := K \cap (C_n \cup \Phi^{-1}(C_n))$$

We already know that

$$\text{Capt}_{\mathcal{S}_\Phi}(K, C) \subset \bigcup_{n \geq 0} C_n$$

Proposition 6.8 *The discrete viable-capture basin*

$$\text{Capt}_{\mathcal{S}_\Phi}(K, C) = \bigcup_{n \geq 0} C_n = \mathcal{B}_{\Phi^{-1}}^b(K, C)$$

is the union of these subsets C_j .

Proof — Let us prove first that C_j is the subset of elements $x \in K$ from which starts a solution $\{x_n\}_{n \geq 0}$ to $x_{n+1} \in \Phi(x_n)$ such that $x_n \in K$ for $n = 0, \dots, j-1$ and $x_j \in C$.

This is obvious for $j = 1$. By induction, assume that it is true for j and let us prove that it is true for $j + 1$: Let x belong to $C_{j+1} := K \cap \Phi^{-1}(C_j)$. There exist $y \in \Phi(x) \cap C_j$, and thus, a sequence $y_{n+1} \in \Phi(y_n)$ such that $y_0 = y$, $y_n \in K$ for $j = 0, \dots, j-1$ and $y_j \in C$. Setting $x_0 := x$ and $x_n := y_{n-1}$ for $n \geq 1$, we see that $x_0 := x$ belongs to K , that $x_1 := y \in \Phi(x) = \Phi(x_0)$ belongs to K , that $x_n \in \Phi(x_{n-1})$ belongs to K for $n \leq j$ and that $x_{j+1} := y_j \in \Phi(x_j)$ belongs to C .

Therefore, if x belongs to one of the subsets C_j , a solution $\{x_n\}_{n \geq 0}$ to the discrete problem $x_{n+1} \in \Phi(x_n)$ is viable in K until it reaches C in j steps. \square

For the absorption basins, Starting with $B_0 := C$, we define recursively the subsets B_n by

$$\forall n \geq 0, B_{n+1} := K \cap (B_n \cup \Phi^{\ominus 1}(B_n))$$

We already know that

$$\text{Abs}_{\mathcal{S}_\Phi}(K, C) \subset \bigcup_{n \geq 0} B_n$$

Proposition 6.9 *If we assume that K and $C \subset K$ are closed and that Φ is lower semicontinuous, then the subsets B_n are closed and*

$$\text{Abs}_{\mathcal{S}_\Phi}(K, C) = \bigcup_{n \geq 0} B_n$$

Proof — Let x belongs to $B_\infty := \bigcup_{n \geq 0} B_n$. If $x \in B_\infty \setminus C$, then for every $n \geq 0$, $x \in \Phi^{\ominus 1}(K_n)$, and thus, $\Phi(x_n) \subset K_n$. Hence

$$\Phi(x) \subset \bigcup_{n \geq 0} B_n =: B_\infty$$

Hence,

$$x \in B_\infty \cap \Phi^{\ominus 1}(B_\infty)$$

and thus, B_∞ is invariant outside C under Φ , and consequently, contained in $\text{Abs}_{\mathcal{S}_\Phi}(K, C)$.

To prove recursively that the subsets B_n are closed, it is enough to prove that $B_{n+1} := B_n \cup \Phi^{\ominus 1}(B_n)$ is closed whenever B_n is closed. Let $x_j \in B_{n+1}$ converge to x , that belongs to B_n . Since Φ is lower semicontinuous, for any $y \in \Phi(x)$, there exists $y_j \in \Phi(x_j)$ converging to y . Since $\Phi(x_j) \subset B_n$, we know that y_j belongs to B_n , and consequently, that y belongs to B_n . Hence $\Phi(x) \subset B_n$, so that x belongs to B_{n+1} . \square

7 Conditional Maps

Let us consider an evolutionary game described by a map $(x, v) \in X \times \mathcal{V} \mapsto \mathcal{S}_v(x) \in \mathcal{C}(0, \infty; X)$, where, for each $v \in \mathcal{V}$, $x \mapsto \mathcal{S}_v(x)$ is an evolutionary system.

In this case, we define the conditional viability kernel as

$$\bigcap_{v \in \mathcal{V}} \text{Viab}_v(K, C)$$

and the guaranteed viability kernel as

$$\bigcup_{v \in \mathcal{V}} \text{Inv}_v(K, C)$$

One can check that only the map $C \mapsto \bigcap_{v \in \mathcal{V}} \text{Viab}_v(K, C)$ is a closing. The map $K \mapsto \bigcap_{v \in \mathcal{V}} \text{Viab}_v(K, C)$ is only a pre-opening, but not necessarily an opening.

However, some of the properties of the conditional and guaranteed viability kernels that we shall review are algebraic in nature.

Let us consider a family of maps $(K, C) \mapsto \mathcal{A}_v(K, C)$ where $v \in \mathcal{V}$ ranges over a subset \mathcal{V} . We associate with \mathcal{A} the conditional map $\mathcal{A}_{\mathcal{V}}$ defined by

$$\mathcal{A}_{\mathcal{V}}(K, C) := \bigcap_{v \in \mathcal{V}} \mathcal{A}_v(K, C)$$

the guaranteed map $\mathcal{A}^{\mathcal{V}}$ defined by

$$\mathcal{A}^{\mathcal{V}}(K, C) := \bigcup_{v \in \mathcal{V}} \mathcal{A}_v(K, C)$$

Lemma 7.1 *If for any $v \in \mathcal{V}$,*

1. *the maps $K \mapsto \mathcal{A}_v(K, C)$ are pre-openings, so are the maps $K \mapsto \mathcal{A}_{\mathcal{V}}(K, C)$ and $K \mapsto \mathcal{A}^{\mathcal{V}}(K, C)$,*
2. *the maps $C \mapsto \mathcal{A}_v(K, C)$ are pre-closings, so are the maps $C \mapsto \mathcal{A}_{\mathcal{V}}(K, C)$ and $K \mapsto \mathcal{A}^{\mathcal{V}}(K, C)$.*

We begin by providing sufficient conditions for the existence of a (unique) minimax (bilateral fixed-point) of the guaranteed map $\mathcal{A}^{\mathcal{V}}$ and the conditional map $\mathcal{A}_{\mathcal{V}}$:

Proposition 7.2 *Let us consider a family of maps $\{\mathcal{A}_v\}_{v \in \mathcal{V}}$ that are pre-openings with respect to K and pre-closings with respect to C .*

1. *If a subset $D \in \mathcal{D}(K, C)$ satisfies*

$$\begin{cases} i) & \forall v \in \mathcal{V}, \mathcal{A}_v(K, D) \subset D \\ ii) & \exists v_0 \in \mathcal{V} \text{ such that } D \subset \mathcal{A}_{v_0}(D, C) \end{cases} \quad (7)$$

then D is equal to $\mathcal{A}^{\mathcal{V}}(K, C)$ and is the unique minimax (bilateral fixed-point) of the map $\mathcal{A}^{\mathcal{V}}$:

$$\mathcal{A}^{\mathcal{V}}(\mathcal{A}^{\mathcal{V}}(K, C), C) = \mathcal{A}^{\mathcal{V}}(K, C) = \mathcal{A}^{\mathcal{V}}(K, \mathcal{A}^{\mathcal{V}}(K, C))$$

2. If a subset $D \in \mathcal{D}(K, C)$ satisfies

$$\begin{cases} i) & \forall v \in \mathcal{V}, D \subset \mathcal{A}_v(D, C) \\ ii) & \exists v_0 \in \mathcal{V} \text{ such that } \mathcal{A}_{v_0}(K, D) \subset D \end{cases} \quad (8)$$

then D is equal to $\mathcal{A}_\mathcal{V}(K, C)$ and is the unique minimax (bilateral fixed-point) of the conditional map $\mathcal{A}_\mathcal{V}$:

$$\mathcal{A}_\mathcal{V}(\mathcal{A}_\mathcal{V}(K, C), C) = \mathcal{A}_\mathcal{V}(K, C) = \mathcal{A}_\mathcal{V}(K, \mathcal{A}_\mathcal{V}(K, C))$$

Proof — By Lemma 7.1, $\mathcal{A}_\mathcal{V}$ is a pre-opening with respect to K and a pre-closing with respect to C .

Condition (7)i) implies that $\mathcal{A}^\mathcal{V}(K, D) \subset D$, and thus, that D belongs to $\mathcal{I}_{\mathcal{A}^\mathcal{V}}(K, C)$. Condition (7)ii) implies that $D \subset \mathcal{A}^{v_0}(K, D) \subset \mathcal{A}^\mathcal{V}(K, D)$, and thus that D belongs to $\mathcal{K}_{\mathcal{A}^\mathcal{V}}(K, C)$. Then $D \in \mathcal{D}(K, C)$ is a minimax (bilateral fixed-point) of $\mathcal{A}^\mathcal{V}$, and thus is equal to $\mathcal{A}^\mathcal{V}(K, C)$ by Proposition 2.3.

The proof of the second statement is the same one. \square

Hence $\mathcal{A}^\mathcal{V}(K, C)$ is the unique candidate for being a of minimax (bilateral fixed-point) of $\mathcal{A}^\mathcal{V}$ and $\mathcal{A}_\mathcal{V}(K, C)$ is the unique candidate for being a of minimax (bilateral fixed-point) of $\mathcal{A}_\mathcal{V}$. Unfortunately, there are examples of maps $\mathcal{A}^\mathcal{V}$ that do not belong to $\mathcal{I}_{\mathcal{A}^\mathcal{V}}(K, C)$, even though the maps \mathcal{A}_v are openings whenever the maps \mathcal{A}_v are openings for all $v \in \mathcal{V}$ and examples of maps $\mathcal{A}_\mathcal{V}$ that do not belong to $\mathcal{K}_{\mathcal{A}_\mathcal{V}}(K, C)$, even though the maps \mathcal{A}_v are closings whenever the maps \mathcal{A}_v are closing for all $v \in \mathcal{V}$:

Lemma 7.3 *Let us assume that for any $v \in \mathcal{V}$, $K \mapsto \mathcal{A}_v(K, C)$ is an opening. Then $K \mapsto \mathcal{A}^\mathcal{V}(K, C)$ is also an opening, so that*

$$\mathcal{A}^\mathcal{V}(\mathcal{A}^\mathcal{V}(K, C), C) = \mathcal{A}^\mathcal{V}(K, C)$$

and

$$\mathcal{A}^\mathcal{V}(K, C) = \bigcup_{D \in \mathcal{K}_{\mathcal{A}^\mathcal{V}}(K, C)} D$$

is the largest fixed point of $D \mapsto \mathcal{A}^\mathcal{V}(D, C)$ between C and K .

If for any $v \in \mathcal{V}$, $C \mapsto \mathcal{A}_v(K, C)$ is a closing, then $C \mapsto \mathcal{A}_\mathcal{V}(K, C)$ is also a closing, so that

$$\mathcal{A}_\mathcal{V}(K, \mathcal{A}_\mathcal{V}(K, C)) = \mathcal{A}_\mathcal{V}(K, C)$$

and

$$\mathcal{A}_\mathcal{V}(K, C) = \bigcap_{D \in \mathcal{I}_{\mathcal{A}_\mathcal{V}}(K, C)} D$$

is the smallest fixed point of $D \mapsto \mathcal{A}_\mathcal{V}(K, D)$ between C and K .

Proof — Indeed, since for any $v \in \mathcal{V}$, the maps $D \mapsto \mathcal{A}_v(D, C)$ are increasing, since $\mathcal{A}_v(K, C) \subset \mathcal{A}^\mathcal{V}(K, C)$, since $\mathcal{A}_v(\mathcal{A}_v(K, C), C) = \mathcal{A}_v(K, C)$ and since $\mathcal{A}_v(D, C) \subset D$, we infer that

$$\left\{ \begin{array}{l} \mathcal{A}^\mathcal{V}(\mathcal{A}^\mathcal{V}(K, C), C) := \bigcup_{v \in \mathcal{V}} \mathcal{A}_v(\mathcal{A}_v(K, C), C) \\ \subset \bigcup_{v \in \mathcal{V}} \mathcal{A}_v(K, C) = \mathcal{A}^\mathcal{V}(K, C) = \bigcup_{v \in \mathcal{V}} \mathcal{A}_v(\mathcal{A}_v(K, C), C) \\ \subset \bigcup_{v \in \mathcal{V}} \mathcal{A}_v(\mathcal{A}_v(K, C), C) \end{array} \right.$$

so that $\mathcal{A}^\mathcal{V}(K, C)$ is the largest fixed point of $D \mapsto \mathcal{A}^\mathcal{V}(D, C)$ between C and K .

In the same way, since for any $v \in \mathcal{V}$, the maps $D \mapsto \mathcal{A}_v(K, D)$ are increasing, since $\mathcal{A}_v(K, C) \subset \mathcal{A}_v(K, D)$, since $\mathcal{A}_v(K, \mathcal{A}_v(K, C)) = \mathcal{A}_v(K, C)$ and since $D \subset \mathcal{A}_v(K, D)$, we infer that

$$\left\{ \begin{array}{l} \mathcal{A}_v(K, \mathcal{A}_v(K, C)) \subset \mathcal{A}_v(K, \mathcal{A}_v(K, C)) \\ := \bigcap_{v \in \mathcal{V}} \mathcal{A}_v(K, \mathcal{A}_v(K, C)) = \bigcap_{v \in \mathcal{V}} \mathcal{A}_v(K, C) = \mathcal{A}_v(K, C) \\ \subset \bigcap_{v \in \mathcal{V}} \mathcal{A}_v(K, \mathcal{A}_v(K, C)) =: \mathcal{A}_v(K, \mathcal{A}_v(K, C)) \end{array} \right.$$

Therefore, $\mathcal{A}_v(K, C)$ is the smallest fixed point of $D \mapsto \mathcal{A}_v(K, D)$ between C and K . \square

The question arises whether we can associate with such a map $\mathcal{A}^\mathcal{V}$ a canonical opening, which in the when $\mathcal{A}_v(K, C) := \text{Inv}_v(K, C)$ case is what is called the discriminating kernel.

They follow from adaptations of the Matheron's theorem.

Indeed, applying the Matheron's Theorem 5.2 to a conditional map $\mathcal{A}_v := \bigcap_{v \in \mathcal{V}} \mathcal{A}_v$, we obtain the following consequence:

Theorem 7.4 *Let us assume that \mathcal{D} is stable for the union and the intersection. If for any $v \in \mathcal{V}$, the maps \mathcal{A}_v are pre-openings with respect to K and extensive with respect to C , and if we assume furthermore that*

1. the map \mathcal{A}_v^\sharp is increasing with respect to C ,
2. the map \mathcal{A}_v^\sharp is idempotent with respect to C :

$$\forall (K, C) \in \mathcal{D}^2, \mathcal{A}_v^\sharp(K, \mathcal{A}_v^\sharp(K, C)) \subset \mathcal{A}_v^\sharp(K, C)$$

then $\mathcal{A}_v^\sharp(K, C)$ is the unique minimax (bilateral fixed-point) $D \in \mathcal{D}(K, C)$ of \mathcal{A}_v^\sharp between C and K :

$$\mathcal{A}_v^\sharp(\mathcal{A}_v^\sharp(K, C), C) = \mathcal{A}_v^\sharp(K, C) = \mathcal{A}_v^\sharp(K, \mathcal{A}_v^\sharp(K, C))$$

8 Discriminating Kernels under Differential Games

8.1 Definition of Evolutionary Games

An evolutionary game is a family of evolutionary systems parametrized by a given family $\tilde{\mathcal{V}}$ of feedbacks $\tilde{v} : x \in X \mapsto \tilde{v}(x) \in \mathcal{V}$.

Definition 8.1 *Let \mathcal{V} be a metric space and $\tilde{\mathcal{V}}$ be a family of feedbacks $\tilde{v} : x \in X \mapsto \tilde{v}(x) \in \mathcal{V}$. Then an evolutionary game is a set-valued map*

$$(x, \tilde{v}) \in X \times \tilde{\mathcal{V}} \rightsquigarrow \mathcal{S}_{\tilde{v}}(x) \in \mathcal{C}(0, \infty; X)$$

such that, for any $\tilde{v} \in \tilde{\mathcal{V}}$, $x \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$ is an evolutionary system.

It is said to be upper semicompact if, for every \tilde{v} , $x \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$ is upper semicompact, and lower semicontinuous if the map $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$ is lower semicontinuous.

Usual differential games provide examples of evolutionary games.

Definition 8.2 *A dynamical game (P, Q, F) is defined by*

- a “set-valued feedback map” $P : X \rightsquigarrow \mathcal{U}$
- a “perturbation” set-valued map $Q : X \rightsquigarrow \mathcal{V}$
- a set-valued map $F : X \times \mathcal{U} \times \mathcal{V} \rightsquigarrow X$ describing the dynamics of the dynamical game:

$$\begin{cases} i) & x'(t) \in F(x(t), u(t), v(t)) \\ ii) & u(t) \in P(x(t)) \\ iii) & v(t) \in Q(x(t)) \end{cases}$$

Naturally, a dynamical game generates the evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$ where $\mathcal{S}_{\tilde{v}}(x)$ is the set of solutions to the differential inclusion

$$x'(t) \in F(x(t), P(x(t)), \tilde{v}(x(t)))$$

starting at x .

8.2 Conditional Viability Kernels and Capture Basins

Definition 8.3 *Let us consider an evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$ and two subsets K and $C \subset K$. We shall say that the subset*

$$\bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C)$$

is the conditional viability kernel with a target under the evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$ and that the subset

$$\bigcap_{\tilde{v}} \text{Capt}_{\tilde{v}}(K, C)$$

is the conditional viable-capture basin of a target under the evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$.

We shall say that K is conditionally viable outside C if

$$K \subset \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C)$$

i.e., if for any state $x_0 \in K$, for all feedbacks \tilde{v} played by Victor, there exists at least one evolution $x(\cdot) \in \mathcal{S}_{\tilde{v}}(x_0)$ viable in K forever or until it reaches C .

We shall say that K traps $C \subset K$ in [27, 31, Cardaliaguet] under the evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$ if

$$K \subset \bigcap_{\tilde{v}} \text{Capt}_{\tilde{v}}(K, C)$$

i.e., if for any state $x_0 \in K$, for all feedbacks \tilde{v} played by Victor, there exists at least one evolution $x(\cdot) \in \mathcal{S}_{\tilde{v}}(x_0)$ viable in K until it reaches C in finite time.

8.3 Fixed-Point Characterization

Proposition 7.2 implies

Proposition 8.4 *If a subset $D \in \mathcal{D}(K, C)$ satisfies*

$$\begin{cases} i) & \forall \tilde{v} \in \tilde{\mathcal{V}}, D \subset \text{Viab}_{\tilde{v}}(K, D) \\ ii) & \exists \tilde{v}_0 \in \tilde{\mathcal{V}} \text{ such that } \text{Viab}_{\tilde{v}_0}(D, C) \subset D \end{cases} \quad (9)$$

then D is equal to $\bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C)$ and is the unique minimax (bilateral fixed-point) of $\bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(\cdot, \cdot)$:

$$\bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}} \left(\bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C), C \right) = \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C) = \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}} \left(K, \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C) \right)$$

We deduce from Theorem 3.3 and Lemma 7.3 the following consequence:

Theorem 8.5 *The map $C \in \mathcal{D}(K, C) \mapsto \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C)$ is a closing so that the conditional viability kernel $\bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C)$ of a subset K with target $C \subset K$ is the smallest subset $D \in \mathcal{D}(K, C)$ satisfying:*

$$\bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C) = \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}} \left(K, \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C) \right)$$

It is antiextensive with respect to K and extensive with respect to C :

$$C \subset \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C) \subset K$$

The same properties hold true for the conditional viable-capture map $\bigcap_{\tilde{v}} \text{Capt}_{\tilde{v}}(K, C)$ of a target C viable in K in a conditional way.

Unfortunately, the conditional viability kernel $\bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C)$ of K outside the target $C \subset K$ is not itself necessarily conditionally viable outside the target C , i.e., it is not a fixed point of the map $K \mapsto \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C)$. In other words, this map is a pre-opening, but not necessarily an opening.

Pierre Cardaliaguet introduced in [27, 31, Cardaliaguet] the opening associated with this map, that he called the discriminating kernel $\text{Disc}(K, C)$ of K with target C :

Definition 8.6 *Let us consider an evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$ and two subsets K and $C \subset K$.*

We shall say that the largest closed subset $D \in \mathcal{D}(K, C)$ conditionally viable in K outside C is the discriminating kernel of K with target C under the evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$, denoted by $\text{Disc}(K, C)$, and that the largest closed subset $D \in \mathcal{D}(K, C)$ trapping C is the trapping basin under the evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$, denoted by $\text{Trap}(K, C)$.

Matheron's Theorem 5.2 implies that the map $K \mapsto \text{Disc}(K, C)$ is the unique opening onto the family of subsets $D \in \mathcal{D}(K, C)$ conditionally viable in K outside C associated with the map $K \in \mathcal{D}(K, C) \mapsto \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C)$ and that $\text{Disc}(K, C)$ is the largest fixed point of this map:

$$\text{Disc}(\text{Disc}(K, C), C) = \text{Disc}(K, C)$$

We shall prove

Proposition 8.7 *Let us consider an evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$.*

Then

$$\begin{cases} i) & C \mapsto \text{Disc}(K, C) \text{ is increasing} \\ ii) & \text{Disc}(K, \text{Disc}(K, C)) = \text{Disc}(K, C) \end{cases} \quad (10)$$

Proof — Let $C_1 \subset C_2 \subset K$ and $x \in \text{Disc}(K, C_1)$. Hence, for any feedback $\tilde{v} \in \tilde{\mathcal{V}}$, there exists at least one evolution $x(\cdot) \in \mathcal{S}_{\tilde{v}}(x)$ viable in $\text{Disc}(K, C_1)$ forever or until it reaches C_1 — and thus, C_2 — in finite time. Hence $\text{Disc}(K, C_1)$ is conditionally viable outside C_2 , so that $\text{Disc}(K, C_1) \subset \text{Disc}(K, C_2)$.

For proving that $\text{Disc}(K, \text{Disc}(K, C))$ traps C , take any $x_0 \in \text{Disc}(K, \text{Disc}(K, C))$. For any \tilde{v} , there exists an evolution $x(\cdot) \in \mathcal{S}_{\tilde{v}}(x_0)$ viable in $\text{Disc}(K, \text{Disc}(K, C))$ forever or else, until it possibly reaches the subset $\text{Disc}(K, C)$ of $\text{Disc}(K, \text{Disc}(K, C))$ at some *finite time* $T > 0$ at $x(T) \in \text{Disc}(K, C)$. In this case, we concatenate $x(\cdot)$ with one solution $y(\cdot) \in \mathcal{S}_{\tilde{v}}(x(T))$ starting from $x(T)$ viable in $\text{Disc}(K, C)$, and thus, in $\text{Disc}(K, \text{Disc}(K, C))$, until it possibly reaches C . Hence, for any \tilde{v} , $\text{Disc}(K, \text{Disc}(K, C))$ captures C , i.e., $\text{Disc}(K, \text{Disc}(K, C))$ traps C by the evolutionary game. \square

Theorem 7.4 and Proposition 8.7 imply

Theorem 8.8 *The discriminating kernel $\text{Disc}(K, C)$ of a subset K with target $C \subset K$ is*

1. *the largest subset $D \in \mathcal{D}(K, C)$ conditionally viable outside the target C ,*
2. *the smallest subset $D \in \mathcal{D}(K, C)$ satisfying $\text{Disc}(K, D) = D$,*
3. *the unique minimax (bilateral fixed-point) $D \in \mathcal{D}(K, C)$ in the sense that*

$$D = \text{Disc}(K, D) = \text{Disc}(D, C)$$

It is antiextensive with respect to K and extensive with respect to C :

$$C \subset \text{Disc}(K, C) \subset K$$

is an opening with respect to K , a closing with respect to C . The same properties hold true for the trapping basin map $\text{Trap}(K, C)$ of a target C viable in K .

8.4 The Cardaliaguet Theorem

Pierre Cardaliaguet has related the concepts of discriminating kernels and conditional viability kernels: Since $K \mapsto \text{Disc}(K, C)$ is the opening associated to the map $K \mapsto \bigcap_{\tilde{v}} \text{Viab}_{\tilde{v}}(K, C)$,

we know that setting

$$K_0 := K \ \& \ \forall i \geq 0, \ K_i := \bigcap_{\tilde{v} \in \tilde{\mathcal{V}}} \text{Viab}_{\tilde{v}}(K_{i-1}, C)$$

then

$$\text{Disc}(K, C) \subset \bigcap_{i=1}^{\infty} K_i$$

Under topological assumptions, we obtain the convergence of the above ‘‘Cardaliaguet algorithm (see [27, Cardaliaguet]):

Theorem 8.9 (Cardaliaguet) *Let us assume that the evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$ is upper semicompact. Then*

$$\text{Disc}(K, C) = \bigcap_{i=1}^{\infty} K_i$$

Proof — We have to check that the intersection $K_{\infty} := \bigcap_{i \geq 0} K_i$ of the closed subsets K_i is conditionally viable outside C , i.e., that for each \tilde{v} , K_{∞} captures C under $\mathcal{S}_{\tilde{v}}$. Indeed, for a fixed \tilde{v} ,

$$C \subset K_{i+1} \subset \text{Viab}_{\tilde{v}}(K_i, C) \subset K_i$$

So, K_{∞} is the intersection of the decreasing sequence of the viability kernels $\text{Viab}_{\tilde{v}}(K_i, C)$ viable under the set-valued map $\mathcal{S}_{\tilde{v}}$. Since $\mathcal{S}_{\tilde{v}}$ is assumed to be upper semicompact, the stability theorem of viability kernels implies that K_{∞} itself is viable outside C under this map $\mathcal{S}_{\tilde{v}}$. Hence, K_{∞} is viable outside C under every set-valued maps $\mathcal{S}_{\tilde{v}}$, and thus, is contained in the discriminating kernel $\text{Disc}(K, C)$. \square

8.5 Guaranteed Viability Kernels and Capture Basins

Definition 8.10 *We shall say that the subset*

$$\bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C)$$

is the guaranteed viability kernel with a target under the evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$ and that the subset

$$\bigcup_{\tilde{v}} \text{Abs}_{\tilde{v}}(K, C)$$

is the guaranteed viable-capture basin of a target under the evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$.

We shall say that K is viable outside C in a guaranteed way if

$$K \subset \bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C)$$

i.e., if for any state $x_0 \in K$, there exists a feedback \tilde{v} played by Victor such that all evolutions $x(\cdot) \in \mathcal{S}_{\tilde{v}}(x_0)$ are viable in K forever or until they reach C .

We shall say that K captures C in a guaranteed way or catches C in K under the evolutionary game $(x, \tilde{v}) \rightsquigarrow \mathcal{S}_{\tilde{v}}(x)$

$$K \subset \bigcup_{\tilde{v}} \text{Abs}_{\tilde{v}}(K, C)$$

i.e., if for any state $x_0 \in K$, there exists a feedback \tilde{v} played by Victor such that all evolutions $x(\cdot) \in \mathcal{S}_{\tilde{v}}(x_0)$ reach C (in finite time) before leaving K .

Naturally, this definition depends upon the choice of the set $\tilde{\mathcal{V}}$ of feedbacks defining the evolutionary game.

8.6 Fixed-Point Characterization

Proposition 7.2 implies

Proposition 8.11 *If a subset $D \in \mathcal{D}(K, C)$ satisfies*

$$\begin{cases} i) & \forall \tilde{v} \in \tilde{\mathcal{V}}, \text{Inv}_{\tilde{v}}(K, D) \subset D \\ ii) & \exists \tilde{v}_0 \in \tilde{\mathcal{V}} \text{ such that } D \subset \text{Inv}_{\tilde{v}_0}(D, C) \end{cases} \quad (11)$$

then D is equal to $\bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C)$ and is the unique minimax (bilateral fixed-point) of $\bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(\cdot, \cdot)$:

$$\bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}} \left(\bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C), C \right) = \bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C) = \bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}} \left(K, \bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C) \right)$$

We deduce from Theorem 3.3 and Lemma 7.3 the following consequence:

Theorem 8.12 *The map $C \in \mathcal{D}(K, C) \mapsto \bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C)$ is an opening so that the guaranteed*

viability kernel $\bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C)$ of a subset K with target $C \subset K$ enjoys the guaranteed viability property

$$\bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}} \left(K, \bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C) \right) = \bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C)$$

and is the largest subset $D \in \mathcal{D}(K, C)$ satisfying:

$$D \subset \bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(D, C)$$

It satisfies

$$C \subset \bigcup_{\tilde{v}} \text{Inv}_{\tilde{v}}(K, C) \subset K$$

The same properties hold true for the guaranteed viable-capture map $\bigcup_{\tilde{v}} \text{Abs}_{\tilde{v}}(K, C)$ of a target C viable in K in a guaranteed way.

References

- [1] AUBIN J.-P. (1987) *Smooth and Heavy Solutions to Control Problems*, in **Nonlinear and Convex Analysis**, Eds. B-L. Lin & Simons S., Proceedings in honor of Ky Fan, Lecture Notes in pure and applied mathematics, June 24-26, 1985,
- [2] AUBIN J.-P. (1991) **Viability Theory** Birkhäuser, Boston, Basel, Berlin
- [3] AUBIN J.-P. (1997) **Dynamic Economic Theory: A Viability Approach**, Springer-Verlag
- [4] AUBIN J.-P. (1999) **Mutational and morphological analysis: tools for shape regulation and morphogenesis**, Birkhäuser
- [5] AUBIN J.-P. (1999) **Impulse Differential Inclusions and Hybrid Systems: A Viability Approach**, Lecture Notes, University of California at Berkeley
- [6] AUBIN J.-P. (2000) *Boundary-Value Problems for Systems of First-Order Partial Differential Inclusions*, NoDEA, 7, 61-84
- [7] AUBIN J.-P. (2000) *Boundary-Value Problems for Systems of Hamilton-Jacobi-Bellman Inclusions with Constraints*, submitted to SIAM J. Control
- [8] AUBIN J.-P. (2000) *Optimal impulse control problems and quasi-variational inequalities thirty years later: A viability approach*, in **Contrôle optimal et EDP: Innovations et Applications**, IOS Press
- [9] AUBIN J.-P. (2001) *The Substratum of Impulse and Hybrid Control Systems*, Proceedings of the HSCC 2001 Conference
- [10] AUBIN J.-P. (2001) *Viability Kernels and Capture Basins of Sets under Differential Inclusions*, SIAM J. Control, 40, 853-881
- [11] AUBIN J.-P. (2002) *Dynamic Core of Fuzzy Dynamical Cooperative Games*, Annals of Dynamic Games, Ninth International Symposium on Dynamical Games and Applications, Adelaide, 2000

- [12] AUBIN J.-P. & DORDAN O. (2001) *Dynamical Qualitative Analysis as Hybrid Systems*, preprint
- [13] AUBIN J.-P. & FRANKOWSKA H. (1990) **Set-Valued Analysis**, Birkhäuser, Boston, Basel, Berlin
- [14] AUBIN J.-P. & FRANKOWSKA H. (1990) *Inclusions aux dérivées partielles gouvernant des contrôles de rétroaction*, Comptes-Rendus de l'Académie des Sciences, Paris, 311, 851-856
- [15] AUBIN J.-P. & FRANKOWSKA H. (1991) *Systèmes hyperboliques d'inclusions aux dérivées partielles*, Comptes-Rendus de l'Académie des Sciences, Paris, 312, 271-276
- [16] AUBIN J.-P. & FRANKOWSKA H. (1992) *Hyperbolic systems of partial differential inclusions*, Annali Scuola Normale di Pisa, 18, 541-562
- [17] AUBIN J.-P. & FRANKOWSKA H. (1995) *Partial differential inclusions governing feedback controls*, J. Convex Analysis, 2, 19-40
- [18] AUBIN J.-P. & FRANKOWSKA H. (1996) *The Viability Kernel Algorithm for Computing Value Functions of Infinite Horizon Optimal Control Problems*, J.Math. Anal. Appl., 201, 555-576
- [19] AUBIN J.-P. & HADDAD G. (submitted) *Detectability through Measurements under Impulse Differential Inclusions*,
- [20] AUBIN J.-P. & HADDAD G. (2001) *Cadenced Runs of Impulse and Hybrid Control Systems*, International Journal Robust and Nonlinear Control
- [21] AUBIN J.-P. & HADDAD G. (2001) *Path-dependent impulse and hybrid control systems*, in **Hybrid Systems: Computation and Control**, 119-132, Di Benedetto & Sangiovanni-Vincentelli Eds, Proceedings of the HSCC 2001 Conference, LNCS 2034, Springer-Verlag
- [22] AUBIN J.-P., LYGEROS J., QUINCAMPOIX. M., SASTRY S. & SEUBE N. (to appear) *Impulse Differential Inclusions: A Viability Approach to Hybrid Systems*,
- [23] AUBIN J.-P., PUJAL D. & SAINT-PIERRE P. (2001) *Dynamic Management of Portfolios with Transaction Costs under Contingent Uncertainty*, preprint
- [24] BACELLI F., COHEN G., OLSDER G. & QUADRAT J.-P. (1992) **Synchronisation and linearity**, Wiley
- [25] CARDALIAGUET P. (1992) *Conditions suffisantes de non vacuité du noyau de viabilité*, Comptes-Rendus de l'Académie des Sciences, Série 1, Paris, 314, 797-800
- [26] CARDALIAGUET P. (1992) *Nonemptiness of the viability kernel of a differential inclusion*, J. of Diff. and Integral Equations
- [27] CARDALIAGUET P. (1994) *Domaines discriminants en jeux différentiels*, Thèse de l'Université de Paris-Dauphine

- [28] CARDALIAGUET P. (1996) *A differential game with two players and one target*, SIAM J. Control and Optimization, 34, 4, 1441-1460.
- [29] CARDALIAGUET P. (1997) *Nonsmooth semi-permeable barriers, Isaacs' equation, and application to a differential game with one target and two players*, Appl. Math. Optim., 36, 125-146
- [30] CARDALIAGUET P. (1997) *On the regularity of Semi-Permeable Surfaces in Control Theory with application to the optimal exit-time problem (Part I and II)*, S.I.A.M. Journal on Opt. Control and Opti., 5, 1638-1671
- [31] CARDALIAGUET P. (2000) **Introduction à la théorie des jeux différentiels**, Lecture Notes, Université Paris-Dauphine
- [32] CARDALIAGUET P., QUINCAMPOIX M. & SAINT-PIERRE P. (1994) *Temps optimaux pour des problèmes avec contraintes et sans contrôlabilité locale* Comptes-Rendus de l'Académie des Sciences, Série 1, Paris, 318, 607-612
- [33] CARDALIAGUET P., QUINCAMPOIX M. & SAINT-PIERRE P. (1995) *Contribution à l'étude des jeux différentiels quantitatifs et qualitatifs avec contrainte sur l'état*, Comptes-Rendus de l'Académie des Sciences, 321, 1543-1548
- [34] CARDALIAGUET P., QUINCAMPOIX M. & SAINT-PIERRE P. (1999) *Set-valued numerical methods for optimal control and differential games*, In **Stochastic and differential games. Theory and numerical methods**, Annals of the International Society of Dynamical Games, 177-247 Birkhäuser
- [35] CRUCK E. (2001) *Problèmes de cible sous contraintes d'état pour des systèmes non linéaires avec sauts d'état*, Comptes-Rendus de l'Académie des Sciences, PARIS,
- [36] CRUCK E. (2001) *Target problems under state constraints for nonlinear controlled impulsive systems*, UBO # 01-2001
- [37] DAVEY B.A. & PRIESTLY H.A. (1990) **Introduction to lattices and order**, Cambridge University Press
- [38] FRANKOWSKA H. (1987) *L'équation d'Hamilton-Jacobi contingente*, Comptes-Rendus de l'Académie des Sciences, PARIS, Série 1, 304, 295-298
- [39] FRANKOWSKA H. (1987) *Optimal trajectories associated to a solution of contingent Hamilton-Jacobi equations*, IEEE, 26th, CDC Conference, Los Angeles, December 9-11
- [40] FRANKOWSKA H. (1989) *Optimal trajectories associated to a solution of contingent Hamilton-Jacobi equations*, Applied Mathematics and Optimization, 19, 291-311
- [41] FRANKOWSKA H. (1989) *Hamilton-Jacobi equation: viscosity solutions and generalized gradients*, J. of Math. Analysis and Appl. 141, 21-26
- [42] FRANKOWSKA H. (1991) *Lower semicontinuous solutions to Hamilton-Jacobi-Bellman equations*, Proceedings of 30th CDC Conference, IEEE, Brighton, December 11-13

- [43] FRANKOWSKA H. (1993) *Lower semicontinuous solutions of Hamilton-Jacobi-Bellman equation*, SIAM J. on Control and Optimization,
- [44] GAUBERT S. (1992) *Théorie des systèmes linéaires dans les diodes*, Thèse Ecole des Mines, Paris
- [45] GIARDINA C. R. & DOUGHERTY E. R. (1988) **Morphological methods in image and signal processing**, Prentice Hall, Englewood Cliffs, NJ.
- [46] GONDRAN M. & MINOUX M. (1984) *Linear algebra in dioids: a survey of recent results*, Annals of Discrete Mathematics, 19, 147-164
- [47] HADDAD G. (1981) *Monotone trajectories of differential inclusions with memory*, Isr. J. Math., 39, 83-100
- [48] HADDAD G. (1981) *Monotone viable trajectories for functional differential inclusions*, J. Diff. Eq., 42, 1-24
- [49] HADDAD G. (1981) *Topological properties of the set of solutions for functional differential differential inclusions*, Nonlinear Anal. Theory, Meth. Appl., 5, 1349-1366
- [50] HEIJMANS H. (1994) **Morphological image operators**, Academic Press
- [51] WINSKEL G. (1993) **The formal semantics of programming languages**, MIT Press
- [52] MATHERON G. (1975) **Random sets and integral geometry**, Wiley
- [53] MATTIOLI J. (1993) *Problèmes inverses et relations différentielles en morphologie mathématique*, Thèse de l' Université de Paris-Dauphine
- [54] MATTIOLI J., DOYEN L. & NAJMAN L. (1996) *Lattice operators underlying dynamic systems*, Set-Valued Analysis
- [55] MATTIOLI J. & SCHMITT M. (1993) **Morphologie mathématique**, Masson
- [56] PUJAL D. (2000) *Valuation et gestion dynamiques de portefeuilles*, Thèse de l'Université de Paris-Dauphine
- [57] QUINCAMPOIX M. (1992) *Enveloppes d'invariance pour des inclusions différentielles Lipschitziennes : applications aux problèmes de cibles*, Comptes-Rendus de l'Académie des Sciences, Paris, 314, 343-347
- [58] QUINCAMPOIX M. (1990) *Frontières de domaines d'invariance et de viabilité pour des inclusions différentielles avec contraintes*, Comptes-Rendus de l'Académie des Sciences, Paris, 311, 411-416
- [59] QUINCAMPOIX M. & SAINT-PIERRE P. (1995) *An algorithm for viability kernels in Hölderian case: Approximation by discrete viability kernels*, J. Math. Syst. Est. and Control, Summary : 115-120

- [60] QUINCAMPOIX M. & VELIOV V. (1998) *Viability with a target: theory and applications*, in **Applications of mathematics in engineering**, 47-54, Heron Press
- [61] SAINT-PIERRE P. (1994) *Approximation of the viability kernel*, Applied Mathematics & Optimisation, 29, 187-209
- [62] SAINT-PIERRE P. (to appear) *Approximation of capture basins for hybrid systems*,
- [63] SERRA J. (1982) **Image Analysis and Mathematical Morphology**, Academic Press
- [64] SERRA J. (1988) **Image Analysis and Mathematical Morphology, Volume 2: Theoretical Advances**, Academic Press
- [65] SHI SHUZHONG (1986) *Théorèmes de viabilité pour les inclusions aux dérivées partielles*, C. R. Acad. Sc. Paris, 303, Série I, 11-14
- [66] SHI SHUZHONG (1988) *Nagumo type condition for partial differential inclusions*, Non-linear Analysis, T.M.A., 12, 951-967
- [67] SHI SHUZHONG (1989) *Viability theorems for a class of differential-operator inclusions*, J. of Diff. Eq., 79, 232-257
- [68] WINSKEL G. (1993) **The formal semantics of programming languages**, MIT Press